



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2012

---

## **Combined search for the quarks of a sequential fourth generation**

CMS Collaboration ; Chatrchyan, S ; Khachatryan, V ; Sirunyan, A M ; Tumasyan, A ; Amsler, C ;  
Chiochia, V ; De Visscher, S ; Favaro, C ; Iova Rikova, M ; Millan Mejias, B ; Otiougova, P ;  
Robmann, P ; Snoek, H ; Tupputi, S ; Verzetti, M

**Abstract:** Results are presented from a search for a fourth generation of quarks produced singly or in pairs in a data set corresponding to an integrated luminosity of 5 fb<sup>-1</sup> recorded by the CMS experiment at the LHC in 2011. A novel strategy has been developed for a combined search for quarks of the up and down type in decay channels with at least one isolated muon or electron. Limits on the mass of the fourth-generation quarks and the relevant Cabibbo-Kobayashi-Maskawa matrix elements are derived in the context of a simple extension of the standard model with a sequential fourth generation of fermions. The existence of mass-degenerate fourth-generation quarks with masses below 685 GeV is excluded at 95% confidence level for minimal off-diagonal mixing between the third- and the fourth-generation quarks. With a mass difference of 25 GeV between the quark masses, the obtained limit on the masses of the fourth-generation quarks shifts by about  $\pm 20$  GeV. These results significantly reduce the allowed parameter space for a fourth generation of fermions.

DOI: <https://doi.org/10.1103/PhysRevD.86.112003>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-76098>

Journal Article

Originally published at:

CMS Collaboration; Chatrchyan, S; Khachatryan, V; Sirunyan, A M; Tumasyan, A; Amsler, C; Chiochia, V; De Visscher, S; Favaro, C; Iova Rikova, M; Millan Mejias, B; Otiougova, P; Robmann, P; Snoek, H; Tupputi, S; Verzetti, M (2012). Combined search for the quarks of a sequential fourth generation. *Physical Review D (Particles, Fields, Gravitation and Cosmology)*, 86(11):112003.

DOI: <https://doi.org/10.1103/PhysRevD.86.112003>

CERN-PH-EP/2012-251  
2012/09/06

CMS-EXO-11-098

# Combined search for the quarks of a sequential fourth generation

The CMS Collaboration\*

## Abstract

Results are presented from a search for a fourth generation of quarks produced singly or in pairs in a data set corresponding to an integrated luminosity of  $5 \text{ fb}^{-1}$  recorded by the CMS experiment at the LHC in 2011. A novel strategy has been developed for a combined search for quarks of the up- and down-type in decay channels with at least one isolated muon or electron. Limits on the mass of the fourth-generation quarks and the relevant CKM matrix elements are derived in the context of a simple extension of the standard model with a sequential fourth generation of fermions. The existence of mass-degenerate fourth-generation quarks with masses below  $685 \text{ GeV}$  is excluded at 95% confidence level for minimal off-diagonal mixing between the third- and the fourth-generation quarks. With a mass difference of  $25 \text{ GeV}$  between the quark masses, the obtained limit on the masses of the fourth-generation quarks shifts by about  $\pm 20 \text{ GeV}$ . This result significantly reduces the allowed parameter space for a fourth generation of fermions.

*Submitted to Physical Review D*

---

\*See Appendix A for the list of collaboration members



# 1 Introduction

The existence of three generations of fermions has been firmly established experimentally [1]. The possibility of a fourth generation of fermions has not been excluded, although it is strongly constrained by precision measurements of electroweak observables. These observables are mainly influenced by the mass differences between the fourth-generation leptons or quarks. In particular, scenarios with a mass difference between the fourth-generation quarks smaller than the mass of the W boson are preferred and even fourth-generation quarks with degenerate masses are allowed [2, 3].

A new generation of fermions requires not only the existence of two additional quarks and two additional leptons, but also an extension of the Cabibbo-Kobayashi-Maskawa (CKM) [4, 5] and Pontecorvo-Maki-Nakagawa-Sakata (PMNS) [6, 7] matrices. New CKM (quark mixing) and PMNS (lepton mixing) matrix elements are constrained by the requirement of consistency with electroweak precision measurements [8].

Previous searches at hadron colliders have considered either pair production *or* single production of *one* of the fourth-generation quarks [9–15]. The most stringent limits exclude the existence of a down-type (up-type) fourth-generation quark with a mass below 611 (570) GeV [14, 15]. These limits on the quark mass values enter a region where the coupling of fourth-generation quarks to the Higgs field becomes large and perturbative calculations for the weak interaction start to fail, assuming the absence of other phenomena beyond the standard model [16]. To increase the sensitivity and to use a consistent approach while searching for a new generation of quarks, we have developed a simultaneous search for the up-type and down-type fourth-generation quarks, based on both the electroweak and strong production mechanisms.

If a fourth generation of quarks exists, their production cross sections and decay branching fractions will be governed by an extended  $4 \times 4$  CKM matrix,  $V_{\text{CKM}}^{4 \times 4}$ , in which we denote the up- and down-type fourth-generation quarks as  $t'$  and  $b'$  respectively. For simplicity, we assume a model with one free parameter,  $A$ , where  $0 \leq A \leq 1$ :

$$\begin{aligned} V_{\text{CKM}}^{4 \times 4} &= \begin{pmatrix} V_{ud} & V_{us} & V_{ub} & V_{ub'} \\ V_{cd} & V_{cs} & V_{cb} & V_{cb'} \\ V_{td} & V_{ts} & V_{tb} & V_{tb'} \\ V_{t'd} & V_{t's} & V_{t'b} & V_{t'b'} \end{pmatrix} \\ &= \begin{pmatrix} \mathcal{O}(1) & \mathcal{O}(0) & \mathcal{O}(0) & 0 \\ \mathcal{O}(0) & \mathcal{O}(1) & \mathcal{O}(0) & 0 \\ \mathcal{O}(0) & \mathcal{O}(0) & \sqrt{A} & \sqrt{1-A} \\ 0 & 0 & -\sqrt{1-A} & \sqrt{A} \end{pmatrix}. \end{aligned}$$

The complex phases are not shown for clarity. Within this model, mixing is allowed only between the third and the fourth generations. This is a reasonable assumption since the mixing between the third and the first two generations is observed to be small [17].

With this search, we set limits on the masses of the fourth-generation quarks as a function of  $A$ . Since  $\sqrt{A} = |V_{tb}|$ , the lower limit of  $|V_{tb}| > 0.81$  from the single-top production cross section measurements [18] translates into a lower limit on the mixing between the third- and fourth-generation quarks in our model of  $A > 0.66$ .

Using the data collected from  $\sqrt{s} = 7$  TeV proton-proton collisions at the Large Hadron Collider (LHC), we search for fourth-generation quarks that are produced in pairs, namely  $b'\bar{b}'$  and  $t'\bar{t}'$ , or through electroweak production, in particular  $tb'$ ,  $t'b$ , and  $t'b'$ , where the charges are omitted in the notation. While the cross sections of the pair production processes do not depend on the

value of  $A$ , the production cross sections of the  $tb'$  and  $t'b$  processes depend linearly on  $(1 - A)$ , and the single-top and  $t'b'$  cross sections on  $A$ .

We assume the  $t'$  and  $b'$  masses to be degenerate within 25 GeV. In the case they are degenerate, they will decay in 100% of the cases to the third-generation quarks, since the decay of one fourth-generation quark to the other is kinematically not allowed. However, even for non-zero mass differences, the branching fractions of the  $t' \rightarrow bW$  and the  $b' \rightarrow tW \rightarrow (bW)W$  decays are close to 100%, provided that the mass difference is small [19]. For instance, for a mass splitting of 25 GeV, and for  $V_{t'b'} = 0.005$  (which would correspond to  $A = 0.99975$  in our model), less than 5% of the decays will be  $b' \rightarrow t'W^*$  (in the case  $m_{t'} < m_{b'}$ ) or  $t' \rightarrow b'W^*$  (in the case  $m_{t'} > m_{b'}$ ). For larger values of  $V_{t'b'}$ , the branching fractions of  $b' \rightarrow t'W^*$  (or  $t' \rightarrow b'W^*$ ) decrease even further. Therefore, the decay chains remain unchanged as long as the mass splitting is relatively small. We expect the following final states:

- $t'b \rightarrow bWb$ ;
- $t'\bar{t}' \rightarrow bWbW$ ;
- $b't \rightarrow tWbW \rightarrow bWWbW$ ;
- $b't' \rightarrow tWbW \rightarrow bWWbW$ ;
- $b'\bar{b}' \rightarrow tWtW \rightarrow bWWbWW$ .

These decay chains imply that two jets from  $b$  quarks and one to four  $W$  bosons are expected in the final state for fourth-generation quarks produced both singly and in pairs. The  $W$  bosons decay to either hadronic or leptonic final states. Events with either one isolated lepton (muon or electron) or two same-sign dileptons or three leptons are selected. The different production processes are classified according to the number of observed  $W$  bosons.

## 2 The Compact Muon Solenoid detector

The central feature of the Compact Muon Solenoid (CMS) detector is a superconducting solenoid, 13 m in length and 6 m in internal diameter, providing an axial magnetic field of 3.8 T. The inside of the solenoid is equipped with various particle detection systems. Charged particle trajectories are measured by a silicon pixel and strip tracker, covering  $0 < \phi < 2\pi$  in azimuth and  $|\eta| < 2.5$ , where the pseudorapidity  $\eta$  is defined as  $-\ln[\tan(\theta/2)]$ , and  $\theta$  is the polar angle of the trajectory with respect to the anticlockwise-beam direction. A crystal electromagnetic calorimeter (ECAL) and a brass/scintillator hadron calorimeter surround the tracking volume and provide high-resolution energy and direction measurements of electrons, photons, and hadronic jets. Muons are measured in gas-ionization detectors embedded in the steel return yoke outside the solenoid. The CMS detector also has extensive forward calorimetry covering up to  $|\eta| < 5$ . The detector is nearly hermetic, allowing for energy balance measurements in the plane transverse to the beam directions. A two-tier trigger system selects the most interesting proton collision events for use in physics analysis. A more detailed description of the CMS detector can be found elsewhere [20].

## 3 Event selection and simulation

The search for the fourth-generation quarks is performed using the  $\sqrt{s} = 7$  TeV proton-proton collisions recorded by the CMS experiment at the LHC. We have analyzed the full dataset collected in 2011 corresponding to an integrated luminosity of  $(5.0 \pm 0.1) \text{ fb}^{-1}$ . Events are selected with a trigger requiring an isolated muon or electron, where the latter is accompanied by at

least one jet identified as a b jet. The muon system, the calorimetry and the tracker are used for the particle-flow (PF) event reconstruction [21]. Jets are reconstructed using the anti- $k_T$  algorithm [22] with a size parameter of 0.5. Events are further selected with at least one high-quality isolated muon or electron with a transverse momentum ( $p_T$ ) exceeding 40 GeV in the acceptance range  $|\eta| < 2.1$  for muons and  $|\eta| < 2.5$  for electrons. The relative isolation,  $I_{\text{rel}}$ , is calculated from the other PF particles within a cone of  $\Delta R = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2} < 0.4$  around the axis of the lepton. It is defined as  $I_{\text{rel}} = (E_T^{\text{charged}} + E_T^{\text{photon}} + E_T^{\text{neutral}})/p_T$ , where  $E_T^{\text{charged}}$  and  $E_T^{\text{photon}}$  are the transverse energies deposited by charged hadrons and photons, respectively, and  $E_T^{\text{neutral}}$  is the transverse energy deposited by neutral particles other than photons. We identify muons and electrons as isolated when  $I_{\text{rel}} < 0.125$  and  $I_{\text{rel}} < 0.1$ , respectively. The requirement on the relative isolation for electrons is tighter than for muons because the backgrounds for electrons are higher than for muons. Electron candidates in the transition region between ECAL barrel and endcap ( $1.44 < |\eta| < 1.57$ ) are excluded because the reconstruction of an electron object in this region is not optimal. We require a missing transverse momentum  $\cancel{E}_T$  of at least 40 GeV. The  $\cancel{E}_T$  is calculated as the absolute value of the vector sum of the  $p_T$  of all reconstructed objects. Jets are required to have a  $p_T > 30$  GeV. The jet energies are corrected to establish a uniform response of the calorimeter in  $\eta$  and a calibrated absolute response in  $p_T$ . Furthermore, a correction is applied to take into account the energy clustered in jets due to additional proton interactions in the same bunch crossing.

The observed data are compared to simulated data generated with POWHEG 301 [23, 24] for the single-top process, PYTHIA 6.4.22 [25] for the diboson processes, and MADGRAPH 5.1.1 [26] for the signal and other standard model processes. The POWHEG and MADGRAPH generators are interfaced with PYTHIA for the decay of the particles as well as the hadronization and the implementation of a CMS custom underlying event tuning (tune Z2) [27]. The matching of the matrix-element partons to the parton showers is obtained using the MLM matching algorithm [28]. The CTEQ6L1 leading-order (LO) parton distributions are used in the event generation [29]. The generated events are passed through the CMS detector simulation based on GEANT4 [30], and then processed by the same reconstruction software as the collision data. The simulated events are reweighted to match the observed distribution of the number of simultaneous proton interactions. For the full dataset collected in 2011, we observe on average about nine interactions in each event. We smear the jet energies in the simulation to match the resolutions measured with data [31]. At least one of the jets within the tracker acceptance ( $|\eta| < 2.4$ ) needs to be identified as a b jet. For the b-jet identification, we require the signed impact parameter significance of the third track in the jet (sorted by decreasing significance) to be larger than a value chosen such that the probability for a light quark jet to be misidentified as a b jet is about 1%. We apply scale factors measured from data to the simulated events to take into account the different b-jet efficiency and the different probability that a light quark or gluon is identified as a b jet in data and simulation [32].

The top-quark pair as well as the W and Z production cross section values used in the analysis correspond to the measured values from CMS [33, 34]. We use the predicted cross section values for the single-top,  $t\bar{t} + W$ ,  $t\bar{t} + Z$ , and same-sign WW processes [35–38]. The cross section values for the diboson production are obtained with the MCFM generator [39, 40].

For the pair-production of the fourth-generation quarks we use the approximate next-to-next-to-leading-order cross section values from [41]. For the electroweak production processes mentioned above, we rescale the next-to-leading-order (NLO) cross sections at 14 TeV [42] to 7 TeV using a scale factor defined as the ratio of the LO cross section at 7 TeV and the LO cross section at 14 TeV as obtained by the MADGRAPH event generator. The resulting production cross

sections are maximal, hence assuming  $|V_{tb'}| = |V_{t'b}| = |V_{t'b'}| = 1$ , and are rescaled according to the value of  $A$ .

## 4 Event classification

Different channels are defined according to the number of  $W$  bosons in the final state. Given that the  $t'$  decay mode is the same as the top-quark decay mode, the  $t'b$  and  $t'\bar{t}'$  processes will yield signatures that are very similar to respectively the single-top and  $t\bar{t}$  processes in the standard model. We select these processes through the single-lepton decay channel. In the signal final states that contain a  $b'$  quark, we expect three or four  $W$  bosons. If two or more of these  $W$  bosons decay to leptons, we may have events with two leptons of the same charge or with three charged leptons. Although the branching fraction of these decays is small compared to that of other decay channels, these final states are very interesting because of the low background that is expected from standard model processes.

### 4.1 The single-electron and single-muon decay channels

On top of the aforementioned event selection criteria, we veto events with additional electrons or muons with  $I_{\text{rel}} < 0.2$  and  $p_T > 10$  GeV for muons and  $p_T > 15$  GeV for electrons. We divide the selected single-lepton events into different subsamples according to the signal final states. Therefore, we define a procedure to count the number of  $W$ -boson candidates. Each event has at least one  $W$  boson that decays to leptons, consistent with the requirements of an isolated lepton and a large missing transverse momentum from the neutrino that escapes detection. The decays of  $W$  bosons to  $q\bar{q}$  final states are reconstructed with the following procedure. For each event, we have a collection of selected jets used as input for the reconstruction of the  $W$ -boson candidates. The one or two jets that are identified as  $b$  jets are removed from the collection.  $W$ -boson candidates are constructed from all possible pairs of the remaining jets in the collection. We use both the expected mass,  $m_W^{\text{fit}} = 84.3$  GeV, and the width,  $\sigma_{m_W}^{\text{fit}} = 9.6$  GeV from a Gaussian fit to the reconstructed mass distribution of jet pairs from the decay of a  $W$  boson in simulated  $t\bar{t}$  events. The  $W$ -boson candidate with a mass that matches the value of  $m_W^{\text{fit}}$  best, is chosen as a  $W$  boson if its mass is within a  $\pm 1\sigma_{m_W}^{\text{fit}}$  window around  $m_W^{\text{fit}}$ . The jet pair that provided the hadronically decaying  $W$  boson is removed from the collection and the procedure is repeated until no more candidates are found for  $W$  bosons decaying to jets. Different exclusive subsamples are defined according to the number of  $b$  jets (exactly one or at least two) and the number of  $W$ -boson candidates (one, two, three, and at least four). There are seven subsamples, because we do not consider the subsample with only one  $b$  jet and one  $W$  boson. The subsample with two  $b$  jets and one  $W$  boson is dominated by singly produced  $t'$  events. In this subsample, we apply a veto for additional jets with a transverse momentum exceeding 30 GeV. Furthermore, since  $b\bar{b}$  background tends to have jets that are produced back-to-back with balanced  $p_T$ , we remove this background by requiring  $\Delta\phi(j_1, j_2) < \frac{\pi}{2} + \pi(|p_T^{j_1} - p_T^{j_2}|) / (p_T^{j_1} + p_T^{j_2})$ .

Table 1 summarizes the requirements that define the different single-lepton decay subsamples, after the criteria on the  $\cancel{E}_T$ , and the lepton and jet  $p_T$  and  $\eta$  are applied.

### 4.2 The same-sign dilepton and trilepton decay channels

The transverse momentum of at least one of the leptons in the multilepton channel is required to be larger than 40 GeV, while the threshold is reduced to 20 GeV for additional leptons. Events with two muons or electrons with a mass within 10 GeV of the  $Z$ -boson mass are rejected to

Table 1: Overview of the event selection requirements defining the different subsamples in the single-lepton decay channel. The single-lepton decay channel is divided in seven different subsamples according to the number of b jets and the number of W-boson candidates.

| single-lepton decay channel        |                               |                            |                            |
|------------------------------------|-------------------------------|----------------------------|----------------------------|
| 1 W                                | 2 W                           | 3 W                        | 4 W                        |
| = 2 jets                           | $\geq 4$ jets                 | $\geq 6$ jets              | $\geq 8$ jets              |
| = 2 b jets                         | either = 1 or $\geq 2$ b jets |                            |                            |
| $\Delta\phi(j_1, j_2)$ requirement | $1 W \rightarrow q\bar{q}$    | $2 W \rightarrow q\bar{q}$ | $3 W \rightarrow q\bar{q}$ |

reduce the standard model background with Z bosons in the final state. We require at least four jets for the same-sign dilepton events. In the case of the trilepton events the minimum number of required jets is reduced to two. Table 2 summarizes the event selection requirements defining the same-sign dilepton and trilepton decay channels that are applied on top of the other requirements on the  $\cancel{E}_T$ , and lepton and jet  $p_T$  and  $\eta$ .

Table 2: Overview of the event selection requirements specific to the same-sign dilepton and trilepton decay channels.

| same-sign dilepton                               | trilepton  |
|--|--|
| = 2 isolated leptons with same sign              | = 3 isolated leptons                             |
| $\geq 4$ jets ( $p_T > 30$ GeV, $ \eta  < 2.4$ ) | $\geq 2$ jets ( $p_T > 30$ GeV, $ \eta  < 2.4$ ) |
| $\geq 1$ b jet                                   | $\geq 1$ b jet                                   |

There are several contributions to the total standard model background for the same-sign dilepton events. One of these contributions comes from events for which the charge of one of the leptons is misreconstructed. Secondly, there are events with one prompt lepton and one non-prompt lepton passing the isolation and identification criteria. Finally, there is an irreducible contribution from standard model processes with two prompt leptons of the same sign; e.g.  $W^\pm W^\pm$ ,  $WZ$ ,  $ZZ$ ,  $t\bar{t} + W$  and  $t\bar{t} + Z$ . Except for  $W^\pm W^\pm$ , these processes are also the main contributions to the total background for the trilepton subsample. The event yields for the irreducible component of the background for the same-sign dilepton channel and the total background in the case of the trilepton subsample are taken from the simulation. We obtain from the data the predicted number of background events for the first two contributions to the total background in the same-sign dilepton subsample.

For the same-sign dilepton events with at least one electron, the background is estimated from control samples. We determine the charge misidentification rate for electrons using a double-isolated-electron trigger. We require two isolated electrons with the dielectron invariant mass within 10 GeV of the Z-boson mass. We select events with  $\cancel{E}_T < 20$  GeV and a transverse mass  $M_T = \sqrt{2p_T^\ell \cancel{E}_T [1 - \cos(\Delta\phi(\ell, \cancel{E}_T))]}$  less than 25 GeV to suppress background from top-quark and W+jets events. We define the charge misidentification ratio  $R$  as the number of events with two electrons of the same sign divided by twice the number of events with two electrons of opposite sign, i.e.  $R = N_{SS}/2N_{OS}$ . We obtain 0.14% and 1.4% for barrel and endcap electron candidates, respectively. After the full event selection is applied, with the exception of the electron sign requirement, we obtain a number of selected data events with two electrons and with an electron and a muon in the final state. The background with two electrons or with an electron and a muon with the same sign is obtained by taking the number of opposite-sign events and scaling it with  $R$ .



Another important background contribution to the same-sign dilepton channel originates from jets being misidentified as an electron or a muon. We require at least one electron or muon with looser isolation ( $I_{\text{rel}} < 0.2$ ) and identification criteria (“loose lepton”). Additionally, we require  $E_T < 20$  GeV and  $M_T < 25$  GeV to suppress background from top-quark and W+jets events. Moreover, we veto events with leptons of the same flavor that have a dilepton mass within 20 GeV of the Z boson mass. We count the number of loose and “tight” (regular isolation and identification criteria) leptons with a  $p_T$  below 35 GeV. The threshold on the  $p_T$  is required to suppress contamination from W+jets events. The probability that a loose (L) lepton passes the tight (T) selection criteria is then given by the ratio  $\epsilon_{TL} = N_T/N_L$ . To estimate the number of events from the background source with a non-prompt lepton, we count the number of events in data that pass the event selection criteria with one lepton passing the tight selection criteria and a second lepton passing the loose but not the tight criteria. This yield is multiplied by  $\epsilon_{TL}(1 - \epsilon_{TL})$  to determine the number of events with a non-prompt lepton in the analysis.

The total number of expected background events for the same-sign dilepton and trilepton channels is given in Table 3.

Table 3: The prediction for the total number of background events compared with the number of observed events in the same-sign dilepton and the trilepton subsamples. The number of expected signal events are also shown for two possible scenarios.

| type                                  | 2 muons         | 2 electrons     | electron+muon   | trilepton       |
|---------------------------------------|-----------------|-----------------|-----------------|-----------------|
| Observed                              | 2               | 2               | 2               | 1               |
| Background                            | $0.83 \pm 0.11$ | $1.36 \pm 0.19$ | $2.27 \pm 0.22$ | $0.96 \pm 0.12$ |
| Signal ( $A = 1, m_{q'} = 550$ GeV)   | $3.31 \pm 0.15$ | $2.03 \pm 0.36$ | $5.29 \pm 0.19$ | $3.37 \pm 0.16$ |
| Signal ( $A = 0.8, m_{q'} = 550$ GeV) | $3.79 \pm 0.15$ | $2.29 \pm 0.36$ | $6.00 \pm 0.19$ | $3.65 \pm 0.16$ |

## 5 Setting lower limits on the fourth-generation quark masses

We have defined different subsamples according to the reconstructed final state. In each of the different subsamples, we reconstruct observables that are sensitive to the presence of the fourth-generation quarks. These observables are used as input to a fit of the combined distributions for the standard model (background-only) hypothesis and the signal-plus-background hypothesis. With the profile likelihood ratio as a test statistic, we calculate the 95% confidence level (CL) upper limits on the combined input cross section of the signal as a function of the  $V_{\text{CKM}}^{4 \times 4}$  parameter  $A$  and the mass of the fourth-generation quarks.

### 5.1 Observables sensitive to the fourth-generation quark production

The expected number of events is small in the subsamples with two leptons of the same sign, the trilepton subsample, and the two single-lepton subsamples with four W-boson candidates. As a consequence, the event counts in each of these subsamples are used as the observable. Table 3 summarizes the event counts for the subsamples with two leptons of the same sign and the trilepton subsample.

In the single-lepton subsamples with one or three W bosons, we use  $S_T$  as the observable to discriminate between the standard model background and the fourth-generation signal, where  $S_T$  is defined as the scalar sum of the transverse momenta of the reconstructed objects in the

final state, namely:

$$S_T = E_T + p_T^\ell + p_T^b + p_T^j + \sum_{i=0}^N p_T^{W_{q\bar{q}}^i}, \quad (1)$$

where the sum runs over the number of reconstructed hadronically decaying W bosons;  $p_T^\ell$  is the  $p_T$  of the lepton,  $p_T^b$  the  $p_T$  of the b jet,  $p_T^j$  the  $p_T$  of the second b jet or, if there is no additional jet identified as a b jet, the  $p_T$  of the jet with the highest transverse momentum in the event that is not used in the W-boson reconstruction, and  $p_T^{W_{q\bar{q}}^i}$  the  $p_T$  of the  $i^{\text{th}}$  reconstructed W boson decaying to jets. In general, the decay products of the fourth-generation quarks are expected to have higher transverse momenta compared to the standard model background. This is shown in Fig. 1 for three of the subsamples. The dominant contribution to the selected signal events in the subsample with two b jets and one W boson would come from the  $t'\bar{b}$  process. Almost no signal events are selected for  $A = 1$ , because in that case the production cross section of  $t'\bar{b}$  is equal to zero. The subsamples with two W bosons are dominated by  $t\bar{t}$  events. In this case we use two sensitive observables,  $S_T$  and the mass of the hadronic bW system,  $m_{bW}$ . The latter observable is sensitive to the fourth generation physics, because of the higher mass of a hypothetical fourth generation  $t'$  quark compared to the top-quark mass. To obtain a higher sensitivity with the  $m_{bW}$  observable, four jets need to be assigned to the quarks to reconstruct the final state  $t'\bar{t}' \rightarrow WbWb \rightarrow q\bar{q}b\ell\nu_\ell b$ . Therefore, six observables with discriminating power between correct and wrong jet/quark assignments are combined with a likelihood ratio method. These observables are angles between the decay products, the W-boson mass, the transverse momentum of the top quark decaying to hadrons, and an observable related to the values of the b-jet identification variable for the jets. The jet/quark assignment with the largest value of the likelihood ratio is chosen. The mass of the bW system is then reconstructed from this chosen jet/quark assignment. The lower plots in Fig. 1 show the projections of the two-dimensional  $S_T$  versus  $m_{bW}$  distribution.

An overview of the observables used in the fit for the presence of the fourth-generation quarks is presented in Table 4.

Table 4: Overview of the observables used in the limit calculation.

| subsample          | observable         |
|--------------------|--------------------|
| single-lepton 1W   | $S_T$              |
| single-lepton 2W   | $S_T$ and $m_{bW}$ |
| single-lepton 3W   | $S_T$              |
| single-lepton 4W   | event yield        |
| same-sign dilepton | event yield        |
| trilepton          | event yield        |

## 5.2 Fitting for the presence of fourth-generation quarks

We construct a single histogram “template” that contains the information of the sensitive observables from all the subsamples. Different template distributions are made for the signal corresponding to the different values of  $A$  and the fourth-generation quark masses  $m_{q'}$ . The binning of the two-dimensional observable distribution in the single-lepton subsamples with two W bosons is defined using the following procedure. We use a binning in the dimension of  $m_{bW}$  such that the top-quark pair background events are uniformly distributed over the bins. Secondly, the binning in the dimension of  $S_T$  in each of the  $m_{bW}$  bins is chosen to obtain uniformly distributed top-quark pair events also in this dimension.




Figure 1: The  $S_T$  distribution for the subsamples with two b jets and one W boson (a), one b jet and three W bosons (b), two b jets and two W bosons (c), and the  $m_{bW}$  distribution for the subsample with two b jets and two W bosons (d). The data distributions of these observables are compared to their expectation from the simulation assuming the fitted nuisance parameters. The fitted values of the nuisance parameters represent the systematic shifts that are applied on the simulation to fit the data in the background-only hypothesis. As an illustration the total uncertainty band is shown around the simulated expected distribution before taking into account the fitted values of the nuisance parameters. The expected distribution for a signal is shown for two different values of the  $V_{CKM}^{4 \times 4}$  parameter  $A$  and for  $b'$  and  $t'$  masses of 550 GeV. The cross section of the signal in the plots is scaled by a factor of eight for visibility. The last bin in all the histograms includes the overflow. We do not expect much signal for  $A = 1$  in (a), because the subsample with two b jets and one W boson is mainly sensitive to single  $t'$  quark production.

The templates of the sensitive observables are used as input to obtain the likelihoods for the background-only and the signal-plus-background hypotheses. Systematic uncertainties are taken into account by introducing nuisance parameters that may affect the shape and the normalization of the templates. In a case where the systematic uncertainty alters the shape of the templates, template morphing [43, 44] is used to interpolate linearly on a bin-by-bin basis between the nominal templates and systematically shifted ones.

The normalization of the templates is affected by the uncertainty in the integrated luminosity, the lepton efficiency and the normalization of the background processes. The integrated luminosity is measured with a precision of 2.2% [45] and has the same normalization effect on all the templates. The uncertainties in the lepton efficiency are a combination of the trigger, selection and identification efficiencies that amount to 3% and 5% for muon and electron respectively. For the uncertainty in the normalization of the background processes, we use the uncertainties in the production cross section of the various standard model processes. The most important contributions that affect the normalization of the templates are the 12% [33] (30%) uncertainty for the top-quark pair (single-top) production cross section and a 50% uncertainty for the  $W$  production cross section because of the large fraction of selected events with jets from heavy flavor quarks. For the multilepton channel, we take into account the uncertainties in the background estimation obtained from the data. We also include the uncertainties in the production cross sections of  $Z$  (5% [34]),  $WW$  (35%),  $WZ$  (42%),  $ZZ$  (27%),  $t\bar{t} + W$  (19%),  $t\bar{t} + Z$  (28%) and  $W^\pm W^\pm$  (49%). The uncertainties in the normalization of diboson and top quark pair production in association with a boson are taken from a comparison of the NLO and the LO predictions.

The largest systematic effects on the shape of the templates originate from the jet energy corrections [31] and the scale factors between data and simulation for the  $b$ -jet efficiency and the probability that a light quark or gluon is identified as a  $b$  jet [32]. These effects are estimated by varying the nominal value by  $\pm 1$  standard deviation. The uncertainty in the jet energy resolution of about 10% has a relatively small effect on the expected limits. The same is true for the uncertainty in the modeling of multiple interactions in the same beam crossing. The latter effect is evaluated by varying the average number of interactions in the simulation by 8%.

The probability density functions of the background-only and the signal-plus-background hypothesis are fitted to the data to fix the nuisance parameters in both models. In the signal-plus-background model, an additional variable, defined as the cross section for the fourth-generation signal obtained by combining the separate search channels, is included. In the combined cross section variable the relative fraction of each fourth-generation signal process is fixed according to the probed model parameters  $(A, m_{q'})$ . Using a Gaussian approximation for the probability density function of the test statistic, we determine the 95% CL expected and observed limits on the combined cross section variable using the  $CL_s$  criterion [46–48]. We exclude the point  $(A, m_{q'})$  at the 95% CL if the upper limit on the combined cross section variable is smaller than its predicted value within the fourth-generation model. The procedure is repeated for each value of  $A$  and  $m_{q'}$ .

### 5.3 Results and discussion

We use the  $CL_s$  procedure to calculate the combined limit for the single muon, single electron, same-sign dilepton and trilepton channels. When the value of the  $V_{\text{CKM}}^{4 \times 4}$  parameter  $A$  approaches unity, the standard model single-top and the  $t'b'$  processes reach their maximal values for the production cross section. When the value of  $A$  decreases, the cross section of these processes decreases linearly with  $A$ . At the same time the expected cross section of the  $t'b$  and  $tb'$  processes increases with  $(1 - A)$  and is equal to zero for  $A = 1$ . Therefore, the  $t'b$

and  $tb'$  processes are expected to enhance the sensitivity for fourth-generation quarks when the parameter  $A$  decreases. This is visible in the upper part of Fig. 2 where both the expected and observed limits on  $m_{q'}$  are more stringent for smaller values of  $A$ . For instance, the limit on the fourth-generation quark masses increases by 70 GeV for  $A = 0.9$  compared to the value of the limit for  $A \sim 1$ . While the  $t'b$  and  $tb'$  processes do not contribute for  $A \sim 1$ , the inclusion of the  $t'b'$  process results in a more stringent limit (a difference of about 30 GeV) compared to when this process is not taken into account.

The existence of fourth-generation quarks with degenerate masses is excluded for all parameter values below the line using the assumed model of the  $V_{\text{CKM}}^{4 \times 4}$  matrix. In particular, fourth-generation quarks with a degenerate mass below 685 GeV are excluded at the 95% CL for a parameter value of  $A \sim 1$ . It is worth noting that no limits can be set for  $A$  exactly equal to unity ( $A = 1$ ), because in this special case the fourth-generation quarks would be stable in the assumed model. The analysis is however valid for values of  $A$  extremely close to unity. The distance between the primary vertex and the decay vertex of the fourth-generation quarks is less than 1 mm for  $1 - A > 2 \times 10^{-14}$ , a number obtained using the LO formula for the decay width of the top quark in which the top-quark mass is replaced with a fourth-generation-quark mass of 600 GeV.

Up to now, the masses of the fourth-generation quarks were assumed to be degenerate. However, if a fourth generation of chiral quarks exists, this is not necessarily the case. Therefore, it is interesting to study how the limit would change for non-degenerate quark masses. If we assume non-degenerate masses, another decay channel for the fourth-generation quarks is possible. Namely, the branching fraction for the decay of  $t'$  ( $b'$ ) into  $b'$  ( $t'$ ) and an off-shell  $W$  boson becomes non-zero. For values of the mass splitting up to about 25 GeV, this branching fraction is small as noted in the introduction. We assume a mass splitting of 25 GeV and unchanged branching fractions for the  $t'$  and  $b'$  decays. The sensitivity of the analysis increases or decreases depending on the specific values of the masses and hence the production cross sections of the fourth-generation quarks. The effect of the mass difference between the fourth-generation quarks on the exclusion limit is shown in the bottom plot of Fig. 2 for a  $V_{\text{CKM}}^{4 \times 4}$  parameter  $A \sim 1$ . For instance in case  $m_{t'} = m_{b'} + 25$  GeV ( $m_{t'} = m_{b'} - 25$  GeV), the limit on  $m_{t'}$  increases about +20 (−20) GeV with respect to the degenerate-mass case. To obtain this limit, we do not take into account the electroweak  $t'b'$  process, which results in more conservative exclusion limits. In particular one observes that quarks with degenerate masses below about 655 GeV are excluded at the 95% CL compared to 685 GeV when the  $t'b'$  process is included.

## 6 Summary

Results from a search for a fourth generation of quarks have been presented. A simple model for a unitary CKM matrix has been defined based on a single parameter  $A = |V_{tb}|^2 = |V_{t'b'}|^2$ . Degenerate masses have been assumed for the fourth-generation quarks, hence  $m_{t'} = m_{b'}$ . The information is combined from different subsamples corresponding to different final states with at least one electron or muon. Observables have been constructed in each of the subsamples and used to differentiate between the standard model background and the processes with fourth-generation quarks. With this strategy the search for singly and pair-produced  $t'$  and  $b'$  quarks has been combined in a coherent way into a single analysis. Model-dependent limits are derived on the mass of the quarks and the  $V_{\text{CKM}}^{4 \times 4}$  matrix element  $A$ . The existence of fourth-generation quarks with masses below 685 GeV is excluded at 95% confidence level for minimal off-diagonal mixing between the third- and the fourth-generation quarks. A non-zero cross section for the single fourth-generation quark production processes, corresponding to a value




Figure 2: Top: Exclusion limit on  $m_{t'} = m_{b'}$  as a function of the  $V_{\text{CKM}}^{4 \times 4}$  parameter  $A$ . The parameter values below the solid line are excluded at 95% CL. The inner (outer) band indicates the 68% (95%) confidence interval around the expected limit. The slope indicates the sensitivity of the analysis to the  $t'b$  and  $tb'$  processes. Bottom: For a  $V_{\text{CKM}}^{4 \times 4}$  parameter value  $A \sim 1$ , the exclusion limit on  $m_{t'}$  versus  $m_{t'} - m_{b'}$  is shown. The exclusion limit is calculated for mass differences up to 25 GeV. The existence of up-type fourth-generation quarks with mass values below the observed limit are excluded at the 95% CL.

of the  $V_{\text{CKM}}^{4 \times 4}$  parameter  $A < 1$  gives rise to a more stringent limit. When a mass difference of 25 GeV is assumed between  $t'$  and  $b'$  quarks, the limit on  $m_{t'}$  shifts by about +20 (−20) GeV for  $m_{t'} = m_{b'} + 25$  GeV ( $m_{t'} = m_{b'} - 25$  GeV). These results significantly reduce the allowed parameter space for a fourth generation of fermions and raise the lower limits on the masses of the fourth generation quarks to the region where nonperturbative effects of the weak interactions are important.

## Acknowledgements

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC machine. We thank the technical and administrative staff at CERN and other CMS institutes, and acknowledge support from BMWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, and FAPESP (Brazil); MES (Bulgaria); CERN; CAS, MoST, and NSFC (China); COLCIENCIAS (Colombia); MSES (Croatia); RPF (Cyprus); MoER, SF0690030s09 and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRT (Greece); OTKA and NKTH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); NRF and WCU (Korea); LAS (Lithuania); CINVESTAV, CONACYT, SEP, and UASLP-FAI (Mexico); MSI (New Zealand); PAEC (Pakistan); MSHE and NSC (Poland); FCT (Portugal); JINR (Armenia, Belarus, Georgia, Ukraine, Uzbekistan); MON, RosAtom, RAS and RFBR (Russia); MSTD (Serbia); SEIDI and CPAN (Spain); Swiss Funding Agencies (Switzerland); NSC (Taipei); ThEP, IPST and NECTEC (Thailand); TUBITAK and TAEK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA). Individuals have received support from the Marie-Curie programme and the European Research Council (European Union); the Leventis Foundation; the A. P. Sloan Foundation; the Alexander von Humboldt Foundation; the Austrian Science Fund (FWF); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the Ministry of Education, Youth and Sports (MEYS) of Czech Republic; the Council of Science and Industrial Research, India; the Compagnia di San Paolo (Torino); and the HOMING PLUS programme of Foundation for Polish Science, cofinanced from European Union, Regional Development Fund.

## References

- [1] Particle Data Group Collaboration, “Number of light neutrino types”, *Eur. Phys. J. C* **15** (2000) 357, doi:10.1007/BF02683445.
- [2] H. Flaecher, M. Goebel, J. Haller et al., “Revisiting the global electroweak fit of the Standard Model and beyond with Gfitter”, *Eur. Phys. J. C* **60** (2009) 543, doi:10.1140/epjc/s10052-009-0966-6, arXiv:0811.0009.
- [3] M. Buchkremer, J.-M. Gerard, and F. Maltoni, “Closing in on a perturbative fourth generation”, *JHEP* **06** (2012) 135, doi:10.1007/JHEP06(2012)135, arXiv:1204.5403.
- [4] N. Cabibbo, “Unitary Symmetry and Leptonic Decays”, *Phys. Rev. Lett.* **10** (1963) 531, doi:10.1103/PhysRevLett.10.531.
- [5] M. Kobayashi and T. Maskawa, “CP Violation in the Renormalizable Theory of Weak Interaction”, *Prog. Theor. Phys.* **49** (1973) 652, doi:10.1143/PTP.49.652.

- [6] B. Pontecorvo, “Mesonium and anti-mesonium”, *Sov. Phys. JETP* **6** (1957) 429.
- [7] Z. Maki, M. Nakagawa, and S. Sakata, “Remarks on the Unified Model of Elementary Particles”, *Prog. Theor. Phys.* **28** (1962) 870, doi:10.1143/PTP.28.870.
- [8] E. Asilar, E. Cavlan, O. Dogangun et al., “Oblique parameters and extra generations via OPUCEM”, *Eur. Phys. J. C* **72** (2012) 1966, doi:10.1140/epjc/s10052-012-1966-5, arXiv:1112.0507.
- [9] ATLAS Collaboration, “Search for down-type fourth generation quarks with the ATLAS Detector in events with one lepton and high transverse momentum hadronically decaying W bosons in  $\sqrt{s} = 7$  TeV pp collisions”, *Phys. Rev. Lett.* **109** (2012) 032001, doi:10.1103/PhysRevLett.109.032001, arXiv:1202.6540.
- [10] ATLAS Collaboration, “Search for pair-produced heavy quarks decaying to Wq in the two-lepton channel at  $\sqrt{s} = 7$  TeV with the ATLAS detector”, *Phys. Rev. D* **86** (2012) 012007, doi:10.1103/PhysRevD.86.012007, arXiv:1202.3389.
- [11] ATLAS Collaboration, “Search for same-sign top-quark production and fourth-generation down-type quarks in pp collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector”, *JHEP* **04** (2012) 069, doi:10.1007/JHEP04(2012)069, arXiv:1202.5520.
- [12] ATLAS Collaboration, “Search for Pair Production of a Heavy Quark Decaying to a W Boson and a b Quark in the Lepton+Jets Channel with the ATLAS Detector”, *Phys. Rev. Lett.* **108** (2012) 261802, doi:10.1103/PhysRevLett.108.261802, arXiv:1202.3076.
- [13] CMS Collaboration, “Search for heavy, top-like quark pair production in the dilepton final state in pp collisions at  $\sqrt{s} = 7$  TeV”, (2012). arXiv:1203.5410. Submitted to Phys. Lett. B.
- [14] CMS Collaboration, “Search for heavy bottom-like quarks in 4.9 inverse femtobarns of pp collisions at  $\sqrt{s} = 7$  TeV”, *JHEP* **05** (2012) 123, doi:10.1007/JHEP05(2012)123, arXiv:1204.1088.
- [15] CMS Collaboration, “Search for the pair production of a top-like  $t'$  quark in events with a lepton and at least four jets in pp collisions at  $\sqrt{s} = 7$  TeV”, (2012). arXiv:1209.0471. Submitted to Phys. Lett. B.
- [16] M. S. Chanowitz, M. A. Furman, and I. Hinchliffe, “Weak interactions of ultra heavy fermions”, *Nucl. Phys. B* **153** (1979) 402, doi:10.1016/0550-3213(79)90606-0.
- [17] Particle Data Group Collaboration, “Review of particle physics”, *Phys. Rev. D* **86** (2012) 010001, doi:10.1103/PhysRevD.86.010001.
- [18] D0 Collaboration, “An improved determination of the width of the top quark”, *Phys. Rev. D* **85** (2012) 091104, doi:10.1103/PhysRevD.85.091104, arXiv:1201.4156.
- [19] Y. Chao, K.-F. Chen, S.-K. Chen et al., “Threshold Effects in the Decay of Heavy  $b'$  and  $t'$  Quarks”, *Phys. Rev. D* **84** (2011) 014029, doi:10.1103/PhysRevD.84.014029, arXiv:1101.0592.
- [20] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **03** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.



- [21] CMS Collaboration, “Commissioning of the Particle-Flow Reconstruction in Minimum-Bias and Jet Events from pp Collisions at 7 TeV”, CMS Physics Analysis Summary CMS-PAS-PFT-10-002, (2010).
- [22] M. Cacciari, G. P. Salam, and G. Soyez, “The anti- $k_t$  jet clustering algorithm”, *JHEP* **04** (2008) 063, doi:10.1088/1126-6708/2008/04/063.
- [23] S. Alioli, P. Nason, C. Oleari et al., “NLO single-top production matched with shower in POWHEG: s- and t-channel contributions”, *JHEP* **09** (2009) 111, doi:10.1088/1126-6708/2009/09/111, arXiv:0907.4076.
- [24] E. Re, “Single-top Wt-channel production matched with parton showers using the POWHEG method”, *Eur. Phys. J. C* **71** (2011) 1547, doi:10.1140/epjc/s10052-011-1547-z, arXiv:1009.2450.
- [25] T. Sjöstrand, S. Mrenna, and P. Z. Skands, “PYTHIA 6.4 physics and manual”, *JHEP* **05** (2006) 026, doi:10.1088/1126-6708/2006/05/026, arXiv:hep-ph/0603175.
- [26] J. Alwall, M. Herquet, F. Maltoni et al., “MadGraph 5: going beyond”, *JHEP* **06** (2011) 128, doi:10.1007/JHEP06(2011)128, arXiv:1106.0522.
- [27] R. Field, “Early LHC Underlying Event Data - Findings and Surprises”, (2010). arXiv:1010.3558.
- [28] M. L. Mangano, M. Moretti, F. Piccinini et al., “Matching matrix elements and shower evolution for top-quark production in hadronic collisions”, *JHEP* **01** (2007) 013, doi:10.1088/1126-6708/2007/01/013, arXiv:hep-ph/0611129.
- [29] J. Pumplin, D. R. Stump, J. Huston et al., “New Generation of Parton Distributions with Uncertainties from Global QCD Analysis”, *JHEP* **07** (2002) 012, doi:10.1088/1126-6708/2002/07/012, arXiv:0201195v3.
- [30] J. Allison et al., “Geant4 developments and applications”, *IEEE Trans. Nucl. Sci.* **53** (2006) 270, doi:10.1109/TNS.2006.869826.
- [31] CMS Collaboration, “Determination of Jet Energy Calibration and Transverse Momentum Resolution in CMS”, *JINST* **06** (2011) 11002, doi:10.1088/1748-0221/6/11/P11002, arXiv:1107.4277.
- [32] CMS Collaboration, “b-Jet Identification in the CMS Experiment”, CMS Physics Analysis Summary CMS-PAS-BTV-11-004, (2011).
- [33] CMS Collaboration, “Measurement of the  $t\bar{t}$  Production Cross Section in pp Collisions at 7 TeV in Lepton + Jets Events Using b-quark Jet Identification”, *Phys. Rev. D* **84** (2011) 092004, doi:10.1103/PhysRevD.84.092004, arXiv:1108.3773.
- [34] CMS Collaboration, “Measurement of the inclusive W and Z production cross sections in pp collisions at  $\sqrt{s} = 7$  TeV with the CMS experiment”, *JHEP* **10** (2011) 132, doi:10.1007/JHEP10(2011)132, arXiv:1107.4789.
- [35] N. Kidonakis, “Next-to-next-to-leading-order collinear and soft gluon corrections for t-channel single top quark production”, *Phys. Rev. D* **83** (2011) 091503, doi:10.1103/PhysRevD.83.091503, arXiv:1103.2792.

- [36] N. Kidonakis, “NNLL resummation for s-channel single top quark production”, *Phys. Rev. D* **81** (2010) 054028, doi:10.1103/PhysRevD.81.054028, arXiv:1001.5034.
- [37] N. Kidonakis, “Two-loop soft anomalous dimensions for single top quark associated production with a W- or H-”, *Phys. Rev. D* **82** (2010) 054018, doi:10.1103/PhysRevD.82.054018, arXiv:1005.4451.
- [38] V. Hirschi, R. Frederix, S. Frixione et al., “Automation of one-loop QCD corrections”, *JHEP* **05** (2011) 044, doi:10.1007/JHEP05(2011)044, arXiv:1103.0621.
- [39] J. M. Campbell and R. K. Ellis, “MCFM for the Tevatron and the LHC”, *Nucl. Phys. Proc. Suppl.* **205-206** (2010) 10, doi:10.1016/j.nuclphysbps.2010.08.011, arXiv:1007.3492.
- [40] J. M. Campbell and R. K. Ellis, “An update on vector boson pair production at hadron colliders”, *Phys. Rev. D* **60** (1999) 113006, doi:10.1103/PhysRevD.60.113006, arXiv:9905386.
- [41] M. Aliev, H. Lacker, U. Langenfeld et al., “HATHOR – HAdronic Top and Heavy quarks crOss section calculatoR”, *Comput. Phys. Commun.* **182** (2011) 1034, doi:10.1016/j.cpc.2010.12.040, arXiv:1007.1327.
- [42] J. M. Campbell, R. Frederix, F. Maltoni et al., “NLO predictions for t-channel production of single top and fourth generation quarks at hadron colliders”, *JHEP* **10** (2009) 042, doi:10.1088/1126-6708/2009/10/042, arXiv:0907.3933.
- [43] A. L. Read, “Linear interpolation of histograms”, *Nucl. Instrum. Meth. A* **425** (1999) 357, doi:10.1016/S0168-9002(98)01347-3.
- [44] J. S. Conway, “Nuisance Parameters in Likelihoods for Multisource Spectra”, in *Proceedings of PHYSTAT 2011 Workshop on Statistical Issues Related to Discovery Claims in Search Experiments and Unfolding*, H. B. Propser and L. Lyons, eds., p. 115. CERN, 2011.
- [45] CMS Collaboration, “Absolute Calibration of the Luminosity Measurement at CMS: Winter 2012 Update”, CMS Physics Analysis Summary CMS-PAS-SMP-12-008, (2012).
- [46] G. Cowan, K. Cranmer, E. Gross et al., “Asymptotic formulae for likelihood-based tests of new physics”, *Eur. Phys. J. C* **71** (2011) 1554, doi:10.1140/epjc/s10052-011-1554-0., arXiv:1007.1727.
- [47] T. Junk, “Confidence level computation for combining searches with small statistics”, *Nucl. Instrum. Meth. A* **434** (1999) 435, doi:10.1016/S0168-9002(99)00498-2, arXiv:hep-ex/9902006.
- [48] A. L. Read, “Presentation of search results: the CLs technique”, *J. Phys. G* **28** (2002) 2693, doi:10.1088/0954-3899/28/10/313.



## A The CMS Collaboration

### **Yerevan Physics Institute, Yerevan, Armenia**

S. Chatrchyan, V. Khachatryan, A.M. Sirunyan, A. Tumasyan

### **Institut für Hochenergiephysik der OeAW, Wien, Austria**

W. Adam, E. Aguilo, T. Bergauer, M. Dragicevic, J. Erö, C. Fabjan<sup>1</sup>, M. Friedl, R. Frühwirth<sup>1</sup>, V.M. Ghete, J. Hammer, N. Hörmann, J. Hrubec, M. Jeitler<sup>1</sup>, W. Kiesenhofer, V. Knünz, M. Krammer<sup>1</sup>, I. Krätschmer, D. Liko, I. Mikulec, M. Pernicka<sup>†</sup>, B. Rahbaran, C. Rohringer, H. Rohringer, R. Schöfbeck, J. Strauss, A. Taurok, W. Waltenberger, G. Walzel, E. Widl, C.-E. Wulz<sup>1</sup>

### **National Centre for Particle and High Energy Physics, Minsk, Belarus**

V. Mossolov, N. Shumeiko, J. Suarez Gonzalez

### **Universiteit Antwerpen, Antwerpen, Belgium**

M. Bansal, S. Bansal, T. Cornelis, E.A. De Wolf, X. Janssen, S. Luyckx, L. Mucibello, S. Ochesanu, B. Roland, R. Rougny, M. Selvaggi, Z. Staykova, H. Van Haevermaet, P. Van Mechelen, N. Van Remortel, A. Van Spillbeeck

### **Vrije Universiteit Brussel, Brussel, Belgium**

F. Blekman, S. Blyweert, J. D'Hondt, R. Gonzalez Suarez, A. Kalogeropoulos, M. Maes, A. Olbrechts, W. Van Doninck, P. Van Mulders, G.P. Van Onsem, I. Villella

### **Université Libre de Bruxelles, Bruxelles, Belgium**

B. Clerbaux, G. De Lentdecker, V. Dero, A.P.R. Gay, T. Hreus, A. Léonard, P.E. Marage, A. Mohammadi, T. Reis, L. Thomas, G. Vander Marcken, C. Vander Velde, P. Vanlaer, J. Wang

### **Ghent University, Ghent, Belgium**

V. Adler, K. Beernaert, A. Cimmino, S. Costantini, G. Garcia, M. Grunewald, B. Klein, J. Lellouch, A. Marinov, J. McCartin, A.A. Ocampo Rios, D. Ryckbosch, N. Strobbe, F. Thyssen, M. Tytgat, P. Verwilligen, S. Walsh, E. Yazgan, N. Zaganidis

### **Université Catholique de Louvain, Louvain-la-Neuve, Belgium**

S. Basegmez, G. Bruno, R. Castello, L. Ceard, C. Delaere, T. du Pree, D. Favart, L. Forthomme, A. Giammanco<sup>2</sup>, J. Hollar, V. Lemaitre, J. Liao, O. Militaru, C. Nuttens, D. Pagano, A. Pin, K. Piotrkowski, N. Schul, J.M. Vizan Garcia

### **Université de Mons, Mons, Belgium**

N. Beliy, T. Caebergs, E. Daubie, G.H. Hammad

### **Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil**

G.A. Alves, M. Correa Martins Junior, D. De Jesus Damiao, T. Martins, M.E. Pol, M.H.G. Souza

### **Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil**

W.L. Aldá Júnior, W. Carvalho, A. Custódio, E.M. Da Costa, C. De Oliveira Martins, S. Fonseca De Souza, D. Matos Figueiredo, L. Mundim, H. Nogima, V. Oguri, W.L. Prado Da Silva, A. Santoro, L. Soares Jorge, A. Sznajder

### **Instituto de Fisica Teorica, Universidade Estadual Paulista, Sao Paulo, Brazil**

T.S. Anjos<sup>3</sup>, C.A. Bernardes<sup>3</sup>, F.A. Dias<sup>4</sup>, T.R. Fernandez Perez Tomei, E.M. Gregores<sup>3</sup>, C. Lagana, F. Marinho, P.G. Mercadante<sup>3</sup>, S.F. Novaes, Sandra S. Padula

### **Institute for Nuclear Research and Nuclear Energy, Sofia, Bulgaria**

V. Genchev<sup>5</sup>, P. Iaydjiev<sup>5</sup>, S. Piperov, M. Rodozov, S. Stoykova, G. Sultanov, V. Tcholakov, R. Trayanov, M. Vutova

**University of Sofia, Sofia, Bulgaria**

A. Dimitrov, R. Hadjiiska, V. Kozhuharov, L. Litov, B. Pavlov, P. Petkov

**Institute of High Energy Physics, Beijing, China**

J.G. Bian, G.M. Chen, H.S. Chen, C.H. Jiang, D. Liang, S. Liang, X. Meng, J. Tao, J. Wang, X. Wang, Z. Wang, H. Xiao, M. Xu, J. Zang, Z. Zhang

**State Key Lab. of Nucl. Phys. and Tech., Peking University, Beijing, China**

C. Asawatangtrakuldee, Y. Ban, S. Guo, Y. Guo, W. Li, S. Liu, Y. Mao, S.J. Qian, H. Teng, D. Wang, L. Zhang, B. Zhu, W. Zou

**Universidad de Los Andes, Bogota, Colombia**

C. Avila, J.P. Gomez, B. Gomez Moreno, A.F. Osorio Oliveros, J.C. Sanabria

**Technical University of Split, Split, Croatia**

N. Godinovic, D. Lelas, R. Plestina<sup>6</sup>, D. Polic, I. Puljak<sup>5</sup>

**University of Split, Split, Croatia**

Z. Antunovic, M. Kovac

**Institute Rudjer Boskovic, Zagreb, Croatia**

V. Brigljevic, S. Duric, K. Kadija, J. Luetic, S. Morovic

**University of Cyprus, Nicosia, Cyprus**

A. Attikis, M. Galanti, G. Mavromanolakis, J. Mousa, C. Nicolaou, F. Ptochos, P.A. Razis

**Charles University, Prague, Czech Republic**

M. Finger, M. Finger Jr.

**Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt**

Y. Assran<sup>7</sup>, S. Elgammal<sup>8</sup>, A. Ellithi Kamel<sup>9</sup>, S. Khalil<sup>8</sup>, M.A. Mahmoud<sup>10</sup>, A. Radi<sup>11,12</sup>

**National Institute of Chemical Physics and Biophysics, Tallinn, Estonia**

M. Kadastik, M. Müntel, M. Raidal, L. Rebane, A. Tiko

**Department of Physics, University of Helsinki, Helsinki, Finland**

P. Eerola, G. Fedi, M. Voutilainen

**Helsinki Institute of Physics, Helsinki, Finland**

J. Härkönen, A. Heikkinen, V. Karimäki, R. Kinnunen, M.J. Kortelainen, T. Lampén, K. Lassila-Perini, S. Lehti, T. Lindén, P. Luukka, T. Mäenpää, T. Peltola, E. Tuominen, J. Tuominiemi, E. Tuovinen, D. Ungaro, L. Wendland

**Lappeenranta University of Technology, Lappeenranta, Finland**

K. Banzuzi, A. Karjalainen, A. Korpela, T. Tuuva

**DSM/IRFU, CEA/Saclay, Gif-sur-Yvette, France**

M. Besancon, S. Choudhury, M. Dejardin, D. Denegri, B. Fabbro, J.L. Faure, F. Ferri, S. Ganjour, A. Givernaud, P. Gras, G. Hamel de Monchenault, P. Jarry, E. Locci, J. Malcles, L. Millischer, A. Nayak, J. Rander, A. Rosowsky, I. Shreyber, M. Titov

**Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France**

S. Baffioni, F. Beaudette, L. Benhabib, L. Bianchini, M. Bluj<sup>13</sup>, C. Broutin, P. Busson, C. Charlot, N. Daci, T. Dahms, L. Dobrzynski, R. Granier de Cassagnac, M. Haguenaue, P. Miné, C. Mironov, I.N. Naranjo, M. Nguyen, C. Ochando, P. Paganini, D. Sabes, R. Salerno, Y. Sirois, C. Veelken, A. Zabi

**Institut Pluridisciplinaire Hubert Curien, Université de Strasbourg, Université de Haute Alsace Mulhouse, CNRS/IN2P3, Strasbourg, France**

J.-L. Agram<sup>14</sup>, J. Andrea, D. Bloch, D. Bodin, J.-M. Brom, M. Cardaci, E.C. Chabert, C. Collard, E. Conte<sup>14</sup>, F. Drouhin<sup>14</sup>, C. Ferro, J.-C. Fontaine<sup>14</sup>, D. Gelé, U. Goerlach, P. Juillot, A.-C. Le Bihan, P. Van Hove

**Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France, Villeurbanne, France**

F. Fassi, D. Mercier

**Université de Lyon, Université Claude Bernard Lyon 1, CNRS-IN2P3, Institut de Physique Nucléaire de Lyon, Villeurbanne, France**

S. Beauceron, N. Beaupere, O. Bondu, G. Boudoul, J. Chasserat, R. Chierici<sup>5</sup>, D. Contardo, P. Depasse, H. El Mamouni, J. Fay, S. Gascon, M. Gouzevitch, B. Ille, T. Kurca, M. Lethuillier, L. Mirabito, S. Perries, V. Sordini, Y. Tschudi, P. Verdier, S. Viret

**Institute of High Energy Physics and Informatization, Tbilisi State University, Tbilisi, Georgia**

Z. Tsamalaidze<sup>15</sup>

**RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany**

G. Anagnostou, S. Beranek, M. Edelhoff, L. Feld, N. Heracleous, O. Hindrichs, R. Jussen, K. Klein, J. Merz, A. Ostapchuk, A. Perieanu, F. Raupach, J. Sammet, S. Schael, D. Sprenger, H. Weber, B. Wittmer, V. Zhukov<sup>16</sup>

**RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany**

M. Ata, J. Caudron, E. Dietz-Laursonn, D. Duchardt, M. Erdmann, R. Fischer, A. Güth, T. Hebbeker, C. Heidemann, K. Hoepfner, D. Klingebiel, P. Kreuzer, C. Magass, M. Merschmeyer, A. Meyer, M. Olschewski, P. Papacz, H. Pieta, H. Reithler, S.A. Schmitz, L. Sonnenschein, J. Steggemann, D. Teyssier, M. Weber

**RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany**

M. Bontenackels, V. Cherepanov, Y. Erdogan, G. Flügge, H. Geenen, M. Geisler, W. Haj Ahmad, F. Hoehle, B. Kargoll, T. Kress, Y. Kuessel, A. Nowack, L. Perchalla, O. Pooth, P. Sauerland, A. Stahl

**Deutsches Elektronen-Synchrotron, Hamburg, Germany**

M. Aldaya Martin, J. Behr, W. Behrenhoff, U. Behrens, M. Bergholz<sup>17</sup>, A. Bethani, K. Borras, A. Burgmeier, A. Cakir, L. Calligaris, A. Campbell, E. Castro, F. Costanza, D. Dammann, C. Diez Pardos, G. Eckerlin, D. Eckstein, G. Flucke, A. Geiser, I. Glushkov, P. Gunnellini, S. Habib, J. Hauk, G. Hellwig, H. Jung, M. Kasemann, P. Katsas, C. Kleinwort, H. Kluge, A. Knutsson, M. Krämer, D. Krücker, E. Kuznetsova, W. Lange, W. Lohmann<sup>17</sup>, B. Lutz, R. Mankel, I. Marfin, M. Marienfeld, I.-A. Melzer-Pellmann, A.B. Meyer, J. Mnich, A. Mussgiller, S. Naumann-Emme, J. Olzem, H. Perrey, A. Petrukhin, D. Pitzl, A. Raspereza, P.M. Ribeiro Cipriano, C. Riedl, E. Ron, M. Rosin, J. Salfeld-Nebgen, R. Schmidt<sup>17</sup>, T. Schoerner-Sadenius, N. Sen, A. Spiridonov, M. Stein, R. Walsh, C. Wissing

**University of Hamburg, Hamburg, Germany**

C. Autermann, V. Blobel, J. Draeger, H. Enderle, J. Erfle, U. Gebbert, M. Görner, T. Hermanns, R.S. Høing, K. Kaschube, G. Kaussen, H. Kirschenmann, R. Klanner, J. Lange, B. Mura, F. Nowak, T. Peiffer, N. Pietsch, D. Rathjens, C. Sander, H. Schettler, P. Schleper, E. Schlieckau, A. Schmidt, M. Schröder, T. Schum, M. Seidel, V. Sola, H. Stadie, G. Steinbrück, J. Thomsen, L. Vanelderen

**Institut für Experimentelle Kernphysik, Karlsruhe, Germany**

C. Barth, J. Berger, C. Böser, T. Chwalek, W. De Boer, A. Descroix, A. Dierlamm, M. Feindt, M. Guthoff<sup>5</sup>, C. Hackstein, F. Hartmann, T. Hauth<sup>5</sup>, M. Heinrich, H. Held, K.H. Hoffmann, S. Honc, I. Katkov<sup>16</sup>, J.R. Komaragiri, P. Lobelle Pardo, D. Martschei, S. Mueller, Th. Müller, M. Niegel, A. Nürnberg, O. Oberst, A. Oehler, J. Ott, G. Quast, K. Rabbertz, F. Ratnikov, N. Ratnikova, S. Röcker, A. Scheurer, F.-P. Schilling, G. Schott, H.J. Simonis, F.M. Stober, D. Troendle, R. Ulrich, J. Wagner-Kuhr, S. Wayand, T. Weiler, M. Zeise

**Institute of Nuclear Physics "Demokritos", Aghia Paraskevi, Greece**

G. Daskalakis, T. Gerasis, S. Kesisoglou, A. Kyriakis, D. Loukas, I. Manolakos, A. Markou, C. Markou, C. Mavrommatis, E. Ntomari

**University of Athens, Athens, Greece**

L. Gouskos, T.J. Mertzimekis, A. Panagiotou, N. Saoulidou

**University of Ioánnina, Ioánnina, Greece**

I. Evangelou, C. Foudas, P. Kokkas, N. Manthos, I. Papadopoulos, V. Patras

**KFKI Research Institute for Particle and Nuclear Physics, Budapest, Hungary**

G. Bencze, C. Hajdu, P. Hidas, D. Horvath<sup>18</sup>, F. Sikler, V. Veszpremi, G. Vesztergombi<sup>19</sup>

**Institute of Nuclear Research ATOMKI, Debrecen, Hungary**

N. Beni, S. Czellar, J. Molnar, J. Palinkas, Z. Szillasi

**University of Debrecen, Debrecen, Hungary**

J. Karancsi, P. Raics, Z.L. Trocsanyi, B. Ujvari

**Panjab University, Chandigarh, India**

S.B. Beri, V. Bhatnagar, N. Dhingra, R. Gupta, M. Kaur, M.Z. Mehta, N. Nishu, L.K. Saini, A. Sharma, J.B. Singh

**University of Delhi, Delhi, India**

Ashok Kumar, Arun Kumar, S. Ahuja, A. Bhardwaj, B.C. Choudhary, S. Malhotra, M. Naimuddin, K. Ranjan, V. Sharma, R.K. Shivpuri

**Saha Institute of Nuclear Physics, Kolkata, India**

S. Banerjee, S. Bhattacharya, S. Dutta, B. Gomber, Sa. Jain, Sh. Jain, R. Khurana, S. Sarkar, M. Sharan

**Bhabha Atomic Research Centre, Mumbai, India**

A. Abdulsalam, R.K. Choudhury, D. Dutta, S. Kailas, V. Kumar, P. Mehta, A.K. Mohanty<sup>5</sup>, L.M. Pant, P. Shukla

**Tata Institute of Fundamental Research - EHEP, Mumbai, India**

T. Aziz, S. Ganguly, M. Guchait<sup>20</sup>, M. Maity<sup>21</sup>, G. Majumder, K. Mazumdar, G.B. Mohanty, B. Parida, K. Sudhakar, N. Wickramage

**Tata Institute of Fundamental Research - HECR, Mumbai, India**

S. Banerjee, S. Dugad

**Institute for Research in Fundamental Sciences (IPM), Tehran, Iran**

H. Arfaei, H. Bakhshiansohi<sup>22</sup>, S.M. Etesami<sup>23</sup>, A. Fahim<sup>22</sup>, M. Hashemi, H. Hesari, A. Jafari<sup>22</sup>, M. Khakzad, M. Mohammadi Najafabadi, S. Paktinat Mehdiabadi, B. Safarzadeh<sup>24</sup>, M. Zeinali<sup>23</sup>

**INFN Sezione di Bari <sup>a</sup>, Università di Bari <sup>b</sup>, Politecnico di Bari <sup>c</sup>, Bari, Italy**

M. Abbrescia<sup>a,b</sup>, L. Barbone<sup>a,b</sup>, C. Calabria<sup>a,b,5</sup>, S.S. Chhibra<sup>a,b</sup>, A. Colaleo<sup>a</sup>, D. Creanza<sup>a,c</sup>,

N. De Filippis<sup>a,c,5</sup>, M. De Palma<sup>a,b</sup>, L. Fiore<sup>a</sup>, G. Iaselli<sup>a,c</sup>, L. Lusito<sup>a,b</sup>, G. Maggi<sup>a,c</sup>, M. Maggi<sup>a</sup>, B. Marangelli<sup>a,b</sup>, S. My<sup>a,c</sup>, S. Nuzzo<sup>a,b</sup>, N. Pacifico<sup>a,b</sup>, A. Pompili<sup>a,b</sup>, G. Pugliese<sup>a,c</sup>, G. Selvaggi<sup>a,b</sup>, L. Silvestris<sup>a</sup>, G. Singh<sup>a,b</sup>, R. Venditti, G. Zito<sup>a</sup>

**INFN Sezione di Bologna <sup>a</sup>, Università di Bologna <sup>b</sup>, Bologna, Italy**

G. Abbiendi<sup>a</sup>, A.C. Benvenuti<sup>a</sup>, D. Bonacorsi<sup>a,b</sup>, S. Braibant-Giacomelli<sup>a,b</sup>, L. Brigliadori<sup>a,b</sup>, P. Capiluppi<sup>a,b</sup>, A. Castro<sup>a,b</sup>, F.R. Cavallo<sup>a</sup>, M. Cuffiani<sup>a,b</sup>, G.M. Dallavalle<sup>a</sup>, F. Fabbri<sup>a</sup>, A. Fanfani<sup>a,b</sup>, D. Fasanella<sup>a,b,5</sup>, P. Giacomelli<sup>a</sup>, C. Grandi<sup>a</sup>, L. Guiducci<sup>a,b</sup>, S. Marcellini<sup>a</sup>, G. Masetti<sup>a</sup>, M. Meneghelli<sup>a,b,5</sup>, A. Montanari<sup>a</sup>, F.L. Navarria<sup>a,b</sup>, F. Odorici<sup>a</sup>, A. Perrotta<sup>a</sup>, F. Primavera<sup>a,b</sup>, A.M. Rossi<sup>a,b</sup>, T. Rovelli<sup>a,b</sup>, G.P. Siroli<sup>a,b</sup>, R. Travaglini<sup>a,b</sup>

**INFN Sezione di Catania <sup>a</sup>, Università di Catania <sup>b</sup>, Catania, Italy**

S. Albergo<sup>a,b</sup>, G. Cappello<sup>a,b</sup>, M. Chiorboli<sup>a,b</sup>, S. Costa<sup>a,b</sup>, R. Potenza<sup>a,b</sup>, A. Tricomi<sup>a,b</sup>, C. Tuve<sup>a,b</sup>

**INFN Sezione di Firenze <sup>a</sup>, Università di Firenze <sup>b</sup>, Firenze, Italy**

G. Barbagli<sup>a</sup>, V. Ciulli<sup>a,b</sup>, C. Civinini<sup>a</sup>, R. D'Alessandro<sup>a,b</sup>, E. Focardi<sup>a,b</sup>, S. Frosali<sup>a,b</sup>, E. Gallo<sup>a</sup>, S. Gonzi<sup>a,b</sup>, M. Meschini<sup>a</sup>, S. Paoletti<sup>a</sup>, G. Sguazzoni<sup>a</sup>, A. Tropiano<sup>a</sup>

**INFN Laboratori Nazionali di Frascati, Frascati, Italy**

L. Benussi, S. Bianco, S. Colafranceschi<sup>25</sup>, F. Fabbri, D. Piccolo

**INFN Sezione di Genova <sup>a</sup>, Università di Genova <sup>b</sup>, Genova, Italy**

P. Fabbriatore<sup>a</sup>, R. Musenich<sup>a</sup>, S. Tosi<sup>a,b</sup>

**INFN Sezione di Milano-Bicocca <sup>a</sup>, Università di Milano-Bicocca <sup>b</sup>, Milano, Italy**

A. Benaglia<sup>a,b</sup>, F. De Guio<sup>a,b</sup>, L. Di Matteo<sup>a,b,5</sup>, S. Fiorendi<sup>a,b</sup>, S. Gennai<sup>a,5</sup>, A. Ghezzi<sup>a,b</sup>, S. Malvezzi<sup>a</sup>, R.A. Manzoni<sup>a,b</sup>, A. Martelli<sup>a,b</sup>, A. Massironi<sup>a,b,5</sup>, D. Menasce<sup>a</sup>, L. Moroni<sup>a</sup>, M. Paganoni<sup>a,b</sup>, D. Pedrini<sup>a</sup>, S. Ragazzi<sup>a,b</sup>, N. Redaelli<sup>a</sup>, S. Sala<sup>a</sup>, T. Tabarelli de Fatis<sup>a,b</sup>

**INFN Sezione di Napoli <sup>a</sup>, Università di Napoli "Federico II" <sup>b</sup>, Napoli, Italy**

S. Buontempo<sup>a</sup>, C.A. Carrillo Montoya<sup>a</sup>, N. Cavallo<sup>a,26</sup>, A. De Cosa<sup>a,b,5</sup>, O. Dogangun<sup>a,b</sup>, F. Fabozzi<sup>a,26</sup>, A.O.M. Iorio<sup>a</sup>, L. Lista<sup>a</sup>, S. Meola<sup>a,27</sup>, M. Merola<sup>a,b</sup>, P. Paolucci<sup>a,5</sup>

**INFN Sezione di Padova <sup>a</sup>, Università di Padova <sup>b</sup>, Università di Trento (Trento) <sup>c</sup>, Padova, Italy**

P. Azzi<sup>a</sup>, N. Bacchetta<sup>a,5</sup>, D. Bisello<sup>a,b</sup>, A. Branca<sup>a,b,5</sup>, R. Carlin<sup>a,b</sup>, P. Checchia<sup>a</sup>, T. Dorigo<sup>a</sup>, U. Dosselli<sup>a</sup>, F. Gasparini<sup>a,b</sup>, U. Gasparini<sup>a,b</sup>, A. Gozzelino<sup>a</sup>, K. Kanishchev<sup>a,c</sup>, S. Lacaprara<sup>a</sup>, I. Lazzizzera<sup>a,c</sup>, M. Margoni<sup>a,b</sup>, A.T. Meneguzzo<sup>a,b</sup>, J. Pazzini<sup>a,b</sup>, N. Pozzobon<sup>a,b</sup>, P. Ronchese<sup>a,b</sup>, F. Simonetto<sup>a,b</sup>, E. Torassa<sup>a</sup>, M. Tosi<sup>a,b,5</sup>, S. Vanini<sup>a,b</sup>, P. Zotto<sup>a,b</sup>, G. Zumerle<sup>a,b</sup>

**INFN Sezione di Pavia <sup>a</sup>, Università di Pavia <sup>b</sup>, Pavia, Italy**

M. Gabusi<sup>a,b</sup>, S.P. Ratti<sup>a,b</sup>, C. Riccardi<sup>a,b</sup>, P. Torre<sup>a,b</sup>, P. Vitulo<sup>a,b</sup>

**INFN Sezione di Perugia <sup>a</sup>, Università di Perugia <sup>b</sup>, Perugia, Italy**

M. Biasini<sup>a,b</sup>, G.M. Bilei<sup>a</sup>, L. Fanò<sup>a,b</sup>, P. Lariccia<sup>a,b</sup>, A. Lucaroni<sup>a,b,5</sup>, G. Mantovani<sup>a,b</sup>, M. Menichelli<sup>a</sup>, A. Nappi<sup>a,b,†</sup>, F. Romeo<sup>a,b</sup>, A. Saha<sup>a</sup>, A. Santocchia<sup>a,b</sup>, A. Spiezia<sup>a,b</sup>, S. Taroni<sup>a,b</sup>

**INFN Sezione di Pisa <sup>a</sup>, Università di Pisa <sup>b</sup>, Scuola Normale Superiore di Pisa <sup>c</sup>, Pisa, Italy**

P. Azzurri<sup>a,c</sup>, G. Bagliesi<sup>a</sup>, T. Boccali<sup>a</sup>, G. Broccolo<sup>a,c</sup>, R. Castaldi<sup>a</sup>, R.T. D'Agnolo<sup>a,c</sup>, R. Dell'Orso<sup>a</sup>, F. Fiori<sup>a,b,5</sup>, L. Foà<sup>a,c</sup>, A. Giassi<sup>a</sup>, A. Kraan<sup>a</sup>, F. Ligabue<sup>a,c</sup>, T. Lomtadze<sup>a</sup>, L. Martini<sup>a,28</sup>, A. Messineo<sup>a,b</sup>, F. Palla<sup>a</sup>, A. Rizzi<sup>a,b</sup>, A.T. Serban<sup>a,29</sup>, P. Spagnolo<sup>a</sup>, P. Squillacioti<sup>a,5</sup>, R. Tenchini<sup>a</sup>, G. Tonelli<sup>a,b,5</sup>, A. Venturi<sup>a</sup>, P.G. Verдини<sup>a</sup>

**INFN Sezione di Roma <sup>a</sup>, Università di Roma "La Sapienza" <sup>b</sup>, Roma, Italy**

L. Barone<sup>a,b</sup>, F. Cavallari<sup>a</sup>, D. Del Re<sup>a,b</sup>, M. Diemoz<sup>a</sup>, C. Fanelli, M. Grassi<sup>a,b,5</sup>, E. Longo<sup>a,b</sup>,



P. Meridiani<sup>a,5</sup>, F. Micheli<sup>a,b</sup>, S. Nourbakhsh<sup>a,b</sup>, G. Organtini<sup>a,b</sup>, R. Paramatti<sup>a</sup>, S. Rahatlou<sup>a,b</sup>, M. Sigamani<sup>a</sup>, L. Soffi<sup>a,b</sup>

**INFN Sezione di Torino <sup>a</sup>, Università di Torino <sup>b</sup>, Università del Piemonte Orientale (Novara) <sup>c</sup>, Torino, Italy**

N. Amapane<sup>a,b</sup>, R. Arcidiacono<sup>a,c</sup>, S. Argiro<sup>a,b</sup>, M. Arneodo<sup>a,c</sup>, C. Biino<sup>a</sup>, N. Cartiglia<sup>a</sup>, M. Costa<sup>a,b</sup>, P. De Remigis<sup>a</sup>, N. Demaria<sup>a</sup>, C. Mariotti<sup>a,5</sup>, S. Maselli<sup>a</sup>, E. Migliore<sup>a,b</sup>, V. Monaco<sup>a,b</sup>, M. Musich<sup>a,5</sup>, M.M. Obertino<sup>a,c</sup>, N. Pastrone<sup>a</sup>, M. Pelliccioni<sup>a</sup>, A. Potenza<sup>a,b</sup>, A. Romero<sup>a,b</sup>, R. Sacchi<sup>a,b</sup>, A. Solano<sup>a,b</sup>, A. Staiano<sup>a</sup>, A. Vilela Pereira<sup>a</sup>

**INFN Sezione di Trieste <sup>a</sup>, Università di Trieste <sup>b</sup>, Trieste, Italy**

S. Belforte<sup>a</sup>, V. Candelise<sup>a,b</sup>, F. Cossutti<sup>a</sup>, G. Della Ricca<sup>a,b</sup>, B. Gobbo<sup>a</sup>, M. Marone<sup>a,b,5</sup>, D. Montanino<sup>a,b,5</sup>, A. Penzo<sup>a</sup>, A. Schizzi<sup>a,b</sup>

**Kangwon National University, Chunchon, Korea**

S.G. Heo, T.Y. Kim, S.K. Nam

**Kyungpook National University, Daegu, Korea**

S. Chang, D.H. Kim, G.N. Kim, D.J. Kong, H. Park, S.R. Ro, D.C. Son, T. Son

**Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea**

J.Y. Kim, Zero J. Kim, S. Song

**Korea University, Seoul, Korea**

S. Choi, D. Gyun, B. Hong, M. Jo, H. Kim, T.J. Kim, K.S. Lee, D.H. Moon, S.K. Park

**University of Seoul, Seoul, Korea**

M. Choi, J.H. Kim, C. Park, I.C. Park, S. Park, G. Ryu

**Sungkyunkwan University, Suwon, Korea**

Y. Cho, Y. Choi, Y.K. Choi, J. Goh, M.S. Kim, E. Kwon, B. Lee, J. Lee, S. Lee, H. Seo, I. Yu

**Vilnius University, Vilnius, Lithuania**

M.J. Bilinskas, I. Grigelionis, M. Janulis, A. Juodagalvis

**Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico**

H. Castilla-Valdez, E. De La Cruz-Burelo, I. Heredia-de La Cruz, R. Lopez-Fernandez, R. Magaña Villalba, J. Martínez-Ortega, A. Sánchez-Hernández, L.M. Villasenor-Cendejas

**Universidad Iberoamericana, Mexico City, Mexico**

S. Carrillo Moreno, F. Vazquez Valencia

**Benemerita Universidad Autonoma de Puebla, Puebla, Mexico**

H.A. Salazar Ibarguen

**Universidad Autónoma de San Luis Potosí, San Luis Potosí, Mexico**

E. Casimiro Linares, A. Morelos Pineda, M.A. Reyes-Santos

**University of Auckland, Auckland, New Zealand**

D. Krofcheck

**University of Canterbury, Christchurch, New Zealand**

A.J. Bell, P.H. Butler, R. Doesburg, S. Reucroft, H. Silverwood

**National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan**

M. Ahmad, M.H. Ansari, M.I. Asghar, H.R. Hoorani, S. Khalid, W.A. Khan, T. Khurshid, S. Qazi, M.A. Shah, M. Shoaib

**National Centre for Nuclear Research, Swierk, Poland**

H. Bialkowska, B. Boimska, T. Frueboes, R. Gokieli, M. Górski, M. Kazana, K. Nawrocki, K. Romanowska-Rybinska, M. Szleper, G. Wrochna, P. Zalewski

**Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**

G. Brona, K. Bunkowski, M. Cwiok, W. Dominik, K. Doroba, A. Kalinowski, M. Konecki, J. Krolikowski

**Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal**

N. Almeida, P. Bargassa, A. David, P. Faccioli, P.G. Ferreira Parracho, M. Gallinaro, J. Seixas, J. Varela, P. Vischia

**Joint Institute for Nuclear Research, Dubna, Russia**

I. Belotelov, P. Bunin, M. Gavrilenko, I. Golutvin, A. Kamenev, V. Karjavin, G. Kozlov, A. Lanev, A. Malakhov, P. Moisenz, V. Palichik, V. Perelygin, M. Savina, S. Shmatov, V. Smirnov, A. Volodko, A. Zarubin

**Petersburg Nuclear Physics Institute, Gatchina (St Petersburg), Russia**

S. Evstyukhin, V. Golovtsov, Y. Ivanov, V. Kim, P. Levchenko, V. Murzin, V. Oreshkin, I. Smirnov, V. Sulimov, L. Uvarov, S. Vavilov, A. Vorobyev, An. Vorobyev

**Institute for Nuclear Research, Moscow, Russia**

Yu. Andreev, A. Dermenev, S. Gninenko, N. Golubev, M. Kirsanov, N. Krasnikov, V. Matveev, A. Pashenkov, D. Tlisov, A. Toropin

**Institute for Theoretical and Experimental Physics, Moscow, Russia**

V. Epshteyn, M. Erofeeva, V. Gavrilov, M. Kossov, N. Lychkovskaya, V. Popov, G. Safronov, S. Semenov, V. Stolin, E. Vlasov, A. Zhokin

**Moscow State University, Moscow, Russia**

A. Belyaev, E. Boos, V. Bunichev, M. Dubinin<sup>4</sup>, L. Dudko, A. Gribushin, V. Klyukhin, O. Kodolova, I. Lokhtin, A. Markina, S. Obraztsov, M. Perfilov, S. Petrushanko, A. Popov, L. Sarycheva<sup>†</sup>, V. Savrin, A. Snigirev

**P.N. Lebedev Physical Institute, Moscow, Russia**

V. Andreev, M. Azarkin, I. Dremin, M. Kirakosyan, A. Leonidov, G. Mesyats, S.V. Rusakov, A. Vinogradov

**State Research Center of Russian Federation, Institute for High Energy Physics, Protvino, Russia**

I. Azhgirey, I. Bayshev, S. Bitioukov, V. Grishin<sup>5</sup>, V. Kachanov, D. Konstantinov, A. Korablev, V. Krychkine, V. Petrov, R. Ryutin, A. Sobol, L. Tourtchanovitch, S. Troshin, N. Tyurin, A. Uzunian, A. Volkov

**University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia**

P. Adzic<sup>30</sup>, M. Djordjevic, M. Ekmedzic, D. Krpic<sup>30</sup>, J. Milosevic

**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain**

M. Aguilar-Benitez, J. Alcaraz Maestre, P. Arce, C. Battilana, E. Calvo, M. Cerrada, M. Chamizo

Llatas, N. Colino, B. De La Cruz, A. Delgado Peris, D. Domínguez Vázquez, C. Fernandez Bedoya, J.P. Fernández Ramos, A. Ferrando, J. Flix, M.C. Fouz, P. Garcia-Abia, O. Gonzalez Lopez, S. Goy Lopez, J.M. Hernandez, M.I. Josa, G. Merino, J. Puerta Pelayo, A. Quintario Olmeda, I. Redondo, L. Romero, J. Santaolalla, M.S. Soares, C. Willmott

**Universidad Autónoma de Madrid, Madrid, Spain**

C. Albajar, G. Codispoti, J.F. de Trocóniz

**Universidad de Oviedo, Oviedo, Spain**

H. Brun, J. Cuevas, J. Fernandez Menendez, S. Folgueras, I. Gonzalez Caballero, L. Lloret Iglesias, J. Piedra Gomez

**Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain**

J.A. Brochero Cifuentes, I.J. Cabrillo, A. Calderon, S.H. Chuang, J. Duarte Campderros, M. Felcini<sup>31</sup>, M. Fernandez, G. Gomez, J. Gonzalez Sanchez, A. Graziano, C. Jorda, A. Lopez Virto, J. Marco, R. Marco, C. Martinez Rivero, F. Matorras, F.J. Munoz Sanchez, T. Rodrigo, A.Y. Rodríguez-Marrero, A. Ruiz-Jimeno, L. Scodellaro, M. Sobron Sanudo, I. Vila, R. Vilar Cortabitarte

**CERN, European Organization for Nuclear Research, Geneva, Switzerland**

D. Abbaneo, E. Auffray, G. Auzinger, M. Bachtis, P. Baillon, A.H. Ball, D. Barney, J.F. Benitez, C. Bernet<sup>6</sup>, G. Bianchi, P. Bloch, A. Bocci, A. Bonato, C. Botta, H. Breuker, T. Camporesi, G. Cerminara, T. Christiansen, J.A. Coarasa Perez, D. D'Enterria, A. Dabrowski, A. De Roeck, S. Di Guida, M. Dobson, N. Dupont-Sagorin, A. Elliott-Peisert, B. Frisch, W. Funk, G. Georgiou, M. Giffels, D. Gigi, K. Gill, D. Giordano, M. Giunta, F. Glege, R. Gomez-Reino Garrido, P. Govoni, S. Gowdy, R. Guida, M. Hansen, P. Harris, C. Hartl, J. Harvey, B. Hegner, A. Hinzmann, V. Innocente, P. Janot, K. Kaadze, E. Karavakis, K. Kousouris, P. Lecoq, Y.-J. Lee, P. Lenzi, C. Lourenço, N. Magini, T. Mäki, M. Malberti, L. Malgeri, M. Mannelli, L. Masetti, F. Meijers, S. Mersi, E. Meschi, R. Moser, M.U. Mozer, M. Mulders, P. Musella, E. Nesvold, T. Orimoto, L. Orsini, E. Palencia Cortezon, E. Perez, L. Perrozzi, A. Petrilli, A. Pfeiffer, M. Pierini, M. Pimiä, D. Piparo, G. Polese, L. Quertenmont, A. Racz, W. Reece, J. Rodrigues Antunes, G. Rolandi<sup>32</sup>, C. Rovelli<sup>33</sup>, M. Rovere, H. Sakulin, F. Santanastasio, C. Schäfer, C. Schwick, I. Segoni, S. Sekmen, A. Sharma, P. Siegrist, P. Silva, M. Simon, P. Sphicas<sup>34</sup>, D. Spiga, A. Tsiros, G.I. Veres<sup>19</sup>, J.R. Vlimant, H.K. Wöhri, S.D. Worm<sup>35</sup>, W.D. Zeuner

**Paul Scherrer Institut, Villigen, Switzerland**

W. Bertl, K. Deiters, W. Erdmann, K. Gabathuler, R. Horisberger, Q. Ingram, H.C. Kaestli, S. König, D. Kotlinski, U. Langenegger, F. Meier, D. Renker, T. Rohe, J. Sibille<sup>36</sup>

**Institute for Particle Physics, ETH Zurich, Zurich, Switzerland**

L. Bäni, P. Bortignon, M.A. Buchmann, B. Casal, N. Chanon, A. Deisher, G. Dissertori, M. Dittmar, M. Donegà, M. Dünser, J. Eugster, K. Freudenreich, C. Grab, D. Hits, P. Lecomte, W. Lustermann, A.C. Marini, P. Martinez Ruiz del Arbol, N. Mohr, F. Moortgat, C. Nägeli<sup>37</sup>, P. Nef, F. Nessi-Tedaldi, F. Pandolfi, L. Pape, F. Pauss, M. Peruzzi, F.J. Ronga, M. Rossini, L. Sala, A.K. Sanchez, A. Starodumov<sup>38</sup>, B. Stieger, M. Takahashi, L. Tauscher<sup>†</sup>, A. Thea, K. Theofilatos, D. Treille, C. Urscheler, R. Wallny, H.A. Weber, L. Wehrli

**Universität Zürich, Zurich, Switzerland**

C. Amsler, V. Chiochia, S. De Visscher, C. Favaro, M. Ivova Rikova, B. Millan Mejias, P. Otiougova, P. Robmann, H. Snoek, S. Tuppiti, M. Verzetti

**National Central University, Chung-Li, Taiwan**

Y.H. Chang, K.H. Chen, C.M. Kuo, S.W. Li, W. Lin, Z.K. Liu, Y.J. Lu, D. Mekterovic, A.P. Singh, R. Volpe, S.S. Yu

**National Taiwan University (NTU), Taipei, Taiwan**

P. Bartalini, P. Chang, Y.H. Chang, Y.W. Chang, Y. Chao, K.F. Chen, C. Dietz, U. Grundler, W.-S. Hou, Y. Hsiung, K.Y. Kao, Y.J. Lei, R.-S. Lu, D. Majumder, E. Petrakou, X. Shi, J.G. Shiu, Y.M. Tzeng, X. Wan, M. Wang

**Cukurova University, Adana, Turkey**

A. Adiguzel, M.N. Bakirci<sup>39</sup>, S. Cerci<sup>40</sup>, C. Dozen, I. Dumanoglu, E. Eskut, S. Girgis, G. Gokbulut, E. Gurpinar, I. Hos, E.E. Kangal, T. Karaman, G. Karapinar<sup>41</sup>, A. Kayis Topaksu, G. Onengut, K. Ozdemir, S. Ozturk<sup>42</sup>, A. Polatoz, K. Sogut<sup>43</sup>, D. Sunar Cerci<sup>40</sup>, B. Tali<sup>40</sup>, H. Topakli<sup>39</sup>, L.N. Vergili, M. Vergili

**Middle East Technical University, Physics Department, Ankara, Turkey**

I.V. Akin, T. Aliev, B. Bilin, S. Bilmis, M. Deniz, H. Gamsizkan, A.M. Guler, K. Ocalan, A. Ozpineci, M. Serin, R. Sever, U.E. Surat, M. Yalvac, E. Yildirim, M. Zeyrek

**Bogazici University, Istanbul, Turkey**

E. Gülmez, B. Isildak<sup>44</sup>, M. Kaya<sup>45</sup>, O. Kaya<sup>45</sup>, S. Ozkorucuklu<sup>46</sup>, N. Sonmez<sup>47</sup>

**Istanbul Technical University, Istanbul, Turkey**

K. Cankocak

**National Scientific Center, Kharkov Institute of Physics and Technology, Kharkov, Ukraine**

L. Levchuk

**University of Bristol, Bristol, United Kingdom**

F. Bostock, J.J. Brooke, E. Clement, D. Cussans, H. Flacher, R. Frazier, J. Goldstein, M. Grimes, G.P. Heath, H.F. Heath, L. Kreczko, S. Metson, D.M. Newbold<sup>35</sup>, K. Nirunpong, A. Poll, S. Senkin, V.J. Smith, T. Williams

**Rutherford Appleton Laboratory, Didcot, United Kingdom**

L. Basso<sup>48</sup>, K.W. Bell, A. Belyaev<sup>48</sup>, C. Brew, R.M. Brown, D.J.A. Cockerill, J.A. Coughlan, K. Harder, S. Harper, J. Jackson, B.W. Kennedy, E. Olaiya, D. Petyt, B.C. Radburn-Smith, C.H. Shepherd-Themistocleous, I.R. Tomalin, W.J. Womersley

**Imperial College, London, United Kingdom**

R. Bainbridge, G. Ball, R. Beuselinck, O. Buchmuller, D. Colling, N. Cripps, M. Cutajar, P. Dauncey, G. Davies, M. Della Negra, W. Ferguson, J. Fulcher, D. Futyan, A. Gilbert, A. Guneratne Bryer, G. Hall, Z. Hatherell, J. Hays, G. Iles, M. Jarvis, G. Karapostoli, L. Lyons, A.-M. Magnan, J. Marrouche, B. Mathias, R. Nandi, J. Nash, A. Nikitenko<sup>38</sup>, A. Papageorgiou, J. Pela, M. Pesaresi, K. Petridis, M. Pioppi<sup>49</sup>, D.M. Raymond, S. Rogerson, A. Rose, M.J. Ryan, C. Seez, P. Sharp<sup>†</sup>, A. Sparrow, M. Stoye, A. Tapper, M. Vazquez Acosta, T. Virdee, S. Wakefield, N. Wardle, T. Whyntie

**Brunel University, Uxbridge, United Kingdom**

M. Chadwick, J.E. Cole, P.R. Hobson, A. Khan, P. Kyberd, D. Leggat, D. Leslie, W. Martin, I.D. Reid, P. Symonds, L. Teodorescu, M. Turner

**Baylor University, Waco, USA**

K. Hatakeyama, H. Liu, T. Scarborough

**The University of Alabama, Tuscaloosa, USA**

O. Charaf, C. Henderson, P. Rumerio

**Boston University, Boston, USA**

A. Avetisyan, T. Bose, C. Fantasia, A. Heister, J. St. John, P. Lawson, D. Lazic, J. Rohlf, D. Sperka, L. Sulak

**Brown University, Providence, USA**

J. Alimena, S. Bhattacharya, D. Cutts, A. Ferapontov, U. Heintz, S. Jabeen, G. Kukartsev, E. Laird, G. Landsberg, M. Luk, M. Narain, D. Nguyen, M. Segala, T. Sinthuprasith, T. Speer, K.V. Tsang

**University of California, Davis, Davis, USA**

R. Breedon, G. Breto, M. Calderon De La Barca Sanchez, S. Chauhan, M. Chertok, J. Conway, R. Conway, P.T. Cox, J. Dolen, R. Erbacher, M. Gardner, R. Houtz, W. Ko, A. Kopecky, R. Lander, T. Miceli, D. Pellett, F. Ricci-tam, B. Rutherford, M. Searle, J. Smith, M. Squires, M. Tripathi, R. Vasquez Sierra

**University of California, Los Angeles, Los Angeles, USA**

V. Andreev, D. Cline, R. Cousins, J. Duris, S. Erhan, P. Everaerts, C. Farrell, J. Hauser, M. Ignatenko, C. Jarvis, C. Plager, G. Rakness, P. Schlein<sup>†</sup>, P. Traczyk, V. Valuev, M. Weber

**University of California, Riverside, Riverside, USA**

J. Babb, R. Clare, M.E. Dinardo, J. Ellison, J.W. Gary, F. Giordano, G. Hanson, G.Y. Jeng<sup>50</sup>, H. Liu, O.R. Long, A. Luthra, H. Nguyen, S. Paramesvaran, J. Sturdy, S. Sumowidagdo, R. Wilken, S. Wimpenny

**University of California, San Diego, La Jolla, USA**

W. Andrews, J.G. Branson, G.B. Cerati, S. Cittolin, D. Evans, F. Golf, A. Holzner, R. Kelley, M. Lebourgeois, J. Letts, I. Macneill, B. Mangano, S. Padhi, C. Palmer, G. Petrucciani, M. Pieri, M. Sani, V. Sharma, S. Simon, E. Sudano, M. Tadel, Y. Tu, A. Vartak, S. Wasserbaech<sup>51</sup>, F. Würthwein, A. Yagil, J. Yoo

**University of California, Santa Barbara, Santa Barbara, USA**

D. Barge, R. Bellan, C. Campagnari, M. D'Alfonso, T. Danielson, K. Flowers, P. Geffert, J. Incandela, C. Justus, P. Kalavase, S.A. Koay, D. Kovalskyi, V. Krutelyov, S. Lowette, N. Mccoll, V. Pavlunin, F. Rebassoo, J. Ribnik, J. Richman, R. Rossin, D. Stuart, W. To, C. West

**California Institute of Technology, Pasadena, USA**

A. Apresyan, A. Bornheim, Y. Chen, E. Di Marco, J. Duarte, M. Gataullin, Y. Ma, A. Mott, H.B. Newman, C. Rogan, M. Spiropulu, V. Timciuc, J. Veverka, R. Wilkinson, S. Xie, Y. Yang, R.Y. Zhu

**Carnegie Mellon University, Pittsburgh, USA**

B. Akgun, V. Azzolini, A. Calamba, R. Carroll, T. Ferguson, Y. Iiyama, D.W. Jang, Y.F. Liu, M. Paulini, H. Vogel, I. Vorobiev

**University of Colorado at Boulder, Boulder, USA**

J.P. Cumalat, B.R. Drell, C.J. Edelmaier, W.T. Ford, A. Gaz, B. Heyburn, E. Luigi Lopez, J.G. Smith, K. Stenson, K.A. Ulmer, S.R. Wagner

**Cornell University, Ithaca, USA**

J. Alexander, A. Chatterjee, N. Eggert, L.K. Gibbons, B. Heltsley, A. Khukhunaishvili, B. Kreis, N. Mirman, G. Nicolas Kaufman, J.R. Patterson, A. Ryd, E. Salvati, W. Sun, W.D. Teo, J. Thom, J. Thompson, J. Tucker, J. Vaughan, Y. Weng, L. Winstrom, P. Wittich

**Fairfield University, Fairfield, USA**

D. Winn

**Fermi National Accelerator Laboratory, Batavia, USA**

S. Abdullin, M. Albrow, J. Anderson, L.A.T. Bauerdick, A. Beretvas, J. Berryhill, P.C. Bhat, I. Bloch, K. Burkett, J.N. Butler, V. Chetluru, H.W.K. Cheung, F. Chlebana, V.D. Elvira, I. Fisk, J. Freeman, Y. Gao, D. Green, O. Gutsche, J. Hanlon, R.M. Harris, J. Hirschauer, B. Hooberman, S. Jindariani, M. Johnson, U. Joshi, B. Kilminster, B. Klima, S. Kunori, S. Kwan, C. Leonidopoulos, J. Linacre, D. Lincoln, R. Lipton, J. Lykken, K. Maeshima, J.M. Marraffino, S. Maruyama, D. Mason, P. McBride, K. Mishra, S. Mrenna, Y. Musienko<sup>52</sup>, C. Newman-Holmes, V. O'Dell, O. Prokofyev, E. Sexton-Kennedy, S. Sharma, W.J. Spalding, L. Spiegel, P. Tan, L. Taylor, S. Tkaczyk, N.V. Tran, L. Uplegger, E.W. Vaandering, R. Vidal, J. Whitmore, W. Wu, F. Yang, F. Yumiceva, J.C. Yun

**University of Florida, Gainesville, USA**

D. Acosta, P. Avery, D. Bourilkov, M. Chen, T. Cheng, S. Das, M. De Gruttola, G.P. Di Giovanni, D. Dobur, A. Drozdetskiy, R.D. Field, M. Fisher, Y. Fu, I.K. Furic, J. Gartner, J. Hugon, B. Kim, J. Konigsberg, A. Korytov, A. Kropivnitskaya, T. Kypreos, J.F. Low, K. Matchev, P. Milenovic<sup>53</sup>, G. Mitselmakher, L. Muniz, R. Remington, A. Rinkevicius, P. Sellers, N. Skhirtladze, M. Snowball, J. Yelton, M. Zakaria

**Florida International University, Miami, USA**

V. Gaultney, S. Hewamanage, L.M. Lebolo, S. Linn, P. Markowitz, G. Martinez, J.L. Rodriguez

**Florida State University, Tallahassee, USA**

T. Adams, A. Askew, J. Bochenek, J. Chen, B. Diamond, S.V. Gleyzer, J. Haas, S. Hagopian, V. Hagopian, M. Jenkins, K.F. Johnson, H. Prosper, V. Veeraraghavan, M. Weinberg

**Florida Institute of Technology, Melbourne, USA**

M.M. Baarmand, B. Dorney, M. Hohlmann, H. Kalakhety, I. Vodopiyanov

**University of Illinois at Chicago (UIC), Chicago, USA**

M.R. Adams, I.M. Anghel, L. Apanasevich, Y. Bai, V.E. Bazterra, R.R. Betts, I. Bucinskaite, J. Callner, R. Cavanaugh, O. Evdokimov, L. Gauthier, C.E. Gerber, D.J. Hofman, S. Khalatyan, F. Lacroix, M. Malek, C. O'Brien, C. Silkworth, D. Strom, N. Varelas

**The University of Iowa, Iowa City, USA**

U. Akgun, E.A. Albayrak, B. Bilki<sup>54</sup>, W. Clarida, F. Duru, S. Griffiths, J.-P. Merlo, H. Mermerkaya<sup>55</sup>, A. Mestvirishvili, A. Moeller, J. Nachtman, C.R. Newsom, E. Norbeck, Y. Onel, F. Ozok, S. Sen, E. Tiras, J. Wetzel, T. Yetkin, K. Yi

**Johns Hopkins University, Baltimore, USA**

B.A. Barnett, B. Blumenfeld, S. Bolognesi, D. Fehling, G. Giurgiu, A.V. Gritsan, Z.J. Guo, G. Hu, P. Maksimovic, S. Rappoccio, M. Swartz, A. Whitbeck

**The University of Kansas, Lawrence, USA**

P. Baringer, A. Bean, G. Benelli, O. Grachov, R.P. Kenny Iii, M. Murray, D. Noonan, S. Sanders, R. Stringer, G. Tinti, J.S. Wood, V. Zhukova

**Kansas State University, Manhattan, USA**

A.F. Barfuss, T. Bolton, I. Chakaberia, A. Ivanov, S. Khalil, M. Makouski, Y. Maravin, S. Shrestha, I. Svintradze

**Lawrence Livermore National Laboratory, Livermore, USA**

J. Gronberg, D. Lange, D. Wright

**University of Maryland, College Park, USA**

A. Baden, M. Boutemeur, B. Calvert, S.C. Eno, J.A. Gomez, N.J. Hadley, R.G. Kellogg, M. Kirn,

T. Kolberg, Y. Lu, M. Marionneau, A.C. Mignerey, K. Pedro, A. Peterman, A. Skuja, J. Temple, M.B. Tonjes, S.C. Tonwar, E. Twedt

**Massachusetts Institute of Technology, Cambridge, USA**

A. Apyan, G. Bauer, J. Bendavid, W. Busza, E. Butz, I.A. Cali, M. Chan, V. Dutta, G. Gomez Ceballos, M. Goncharov, K.A. Hahn, Y. Kim, M. Klute, K. Krajczar<sup>56</sup>, W. Li, P.D. Luckey, T. Ma, S. Nahn, C. Paus, D. Ralph, C. Roland, G. Roland, M. Rudolph, G.S.F. Stephans, F. Stöckli, K. Sumorok, K. Sung, D. Velicanu, E.A. Wenger, R. Wolf, B. Wyslouch, M. Yang, Y. Yilmaz, A.S. Yoon, M. Zanetti

**University of Minnesota, Minneapolis, USA**

S.I. Cooper, B. Dahmes, A. De Benedetti, G. Franzoni, A. Gude, S.C. Kao, K. Klapoetke, Y. Kubota, J. Mans, N. Pastika, R. Rusack, M. Sasseville, A. Singovsky, N. Tambe, J. Turkewitz

**University of Mississippi, Oxford, USA**

L.M. Cremaldi, R. Kroeger, L. Perera, R. Rahmat, D.A. Sanders

**University of Nebraska-Lincoln, Lincoln, USA**

E. Avdeeva, K. Bloom, S. Bose, J. Butt, D.R. Claes, A. Dominguez, M. Eads, J. Keller, I. Kravchenko, J. Lazo-Flores, H. Malbouisson, S. Malik, G.R. Snow

**State University of New York at Buffalo, Buffalo, USA**

U. Baur, A. Godshalk, I. Iashvili, S. Jain, A. Kharchilava, A. Kumar, S.P. Shipkowski, K. Smith

**Northeastern University, Boston, USA**

G. Alverson, E. Barberis, D. Baumgartel, M. Chasco, J. Haley, D. Nash, D. Trocino, D. Wood, J. Zhang

**Northwestern University, Evanston, USA**

A. Anastassov, A. Kubik, N. Mucia, N. Odell, R.A. Ofierzynski, B. Pollack, A. Pozdnyakov, M. Schmitt, S. Stoynev, M. Velasco, S. Won

**University of Notre Dame, Notre Dame, USA**

L. Antonelli, D. Berry, A. Brinkerhoff, M. Hildreth, C. Jessop, D.J. Karmgard, J. Kolb, K. Lannon, W. Luo, S. Lynch, N. Marinelli, D.M. Morse, T. Pearson, M. Planer, R. Ruchti, J. Slaunwhite, N. Valls, M. Wayne, M. Wolf

**The Ohio State University, Columbus, USA**

B. Bylsma, L.S. Durkin, C. Hill, R. Hughes, R. Hughes, K. Kotov, T.Y. Ling, D. Puigh, M. Rodenburg, C. Vuosalo, G. Williams, B.L. Winer

**Princeton University, Princeton, USA**

N. Adam, E. Berry, P. Elmer, D. Gerbaudo, V. Halyo, P. Hebda, J. Hegeman, A. Hunt, P. Jindal, D. Lopes Pegna, P. Lujan, D. Marlow, T. Medvedeva, M. Mooney, J. Olsen, P. Piroué, X. Quan, A. Raval, B. Safdi, H. Saka, D. Stickland, C. Tully, J.S. Werner, A. Zuranski

**University of Puerto Rico, Mayaguez, USA**

J.G. Acosta, E. Brownson, X.T. Huang, A. Lopez, H. Mendez, S. Oliveros, J.E. Ramirez Vargas, A. Zatserklyaniy

**Purdue University, West Lafayette, USA**

E. Alagoz, V.E. Barnes, D. Benedetti, G. Bolla, D. Bortoletto, M. De Mattia, A. Everett, Z. Hu, M. Jones, O. Koybasi, M. Kress, A.T. Laasanen, N. Leonardo, V. Maroussov, P. Merkel, D.H. Miller, N. Neumeister, I. Shipsey, D. Silvers, A. Svyatkovskiy, M. Vidal Marono, H.D. Yoo, J. Zablocki, Y. Zheng

**Purdue University Calumet, Hammond, USA**

S. Guragain, N. Parashar

**Rice University, Houston, USA**

A. Adair, C. Boulahouache, K.M. Ecklund, F.J.M. Geurts, B.P. Padley, R. Redjimi, J. Roberts, J. Zabel

**University of Rochester, Rochester, USA**

B. Betchart, A. Bodek, Y.S. Chung, R. Covarelli, P. de Barbaro, R. Demina, Y. Eshaq, A. Garcia-Bellido, P. Goldenzweig, J. Han, A. Harel, D.C. Miner, D. Vishnevskiy, M. Zielinski

**The Rockefeller University, New York, USA**

A. Bhatti, R. Ciesielski, L. Demortier, K. Goulios, G. Lungu, S. Malik, C. Mesropian

**Rutgers, the State University of New Jersey, Piscataway, USA**

S. Arora, A. Barker, J.P. Chou, C. Contreras-Campana, E. Contreras-Campana, D. Duggan, D. Ferencek, Y. Gershtein, R. Gray, E. Halkiadakis, D. Hidas, A. Lath, S. Panwalkar, M. Park, R. Patel, V. Rekovic, J. Robles, K. Rose, S. Salur, S. Schnetzer, C. Seitz, S. Somalwar, R. Stone, S. Thomas

**University of Tennessee, Knoxville, USA**

G. Cerizza, M. Hollingsworth, S. Spanier, Z.C. Yang, A. York

**Texas A&M University, College Station, USA**

R. Eusebi, W. Flanagan, J. Gilmore, T. Kamon<sup>57</sup>, V. Khotilovich, R. Montalvo, I. Osipenkov, Y. Pakhotin, A. Perloff, J. Roe, A. Safonov, T. Sakuma, S. Sengupta, I. Suarez, A. Tatarinov, D. Toback

**Texas Tech University, Lubbock, USA**

N. Akchurin, J. Damgov, C. Dragoiu, P.R. Duder, C. Jeong, K. Kovitanggoon, S.W. Lee, T. Libeiro, Y. Roh, I. Volobouev

**Vanderbilt University, Nashville, USA**

E. Appelt, A.G. Delannoy, C. Florez, S. Greene, A. Gurrola, W. Johns, C. Johnston, P. Kurt, C. Maguire, A. Melo, M. Sharma, P. Sheldon, B. Snook, S. Tuo, J. Velkovska

**University of Virginia, Charlottesville, USA**

M.W. Arenton, M. Balazs, S. Boutle, B. Cox, B. Francis, J. Goodell, R. Hirosky, A. Ledovskoy, C. Lin, C. Neu, J. Wood, R. Yohay

**Wayne State University, Detroit, USA**

S. Gollapinni, R. Harr, P.E. Karchin, C. Kottachchi Kankanamge Don, P. Lamichhane, A. Sakharov

**University of Wisconsin, Madison, USA**

M. Anderson, D. Belknap, L. Borrello, D. Carlsmith, M. Cepeda, S. Dasu, E. Friis, L. Gray, K.S. Grogg, M. Grothe, R. Hall-Wilton, M. Herndon, A. Hervé, P. Klabbers, J. Klukas, A. Lanaro, C. Lazaridis, J. Leonard, R. Loveless, A. Mohapatra, I. Ojalvo, F. Palmonari, G.A. Pierro, I. Ross, A. Savin, W.H. Smith, J. Swanson

†: Deceased

1: Also at Vienna University of Technology, Vienna, Austria

2: Also at National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

3: Also at Universidade Federal do ABC, Santo Andre, Brazil

4: Also at California Institute of Technology, Pasadena, USA



- 5: Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- 6: Also at Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France
- 7: Also at Suez Canal University, Suez, Egypt
- 8: Also at Zewail City of Science and Technology, Zewail, Egypt
- 9: Also at Cairo University, Cairo, Egypt
- 10: Also at Fayoum University, El-Fayoum, Egypt
- 11: Also at British University, Cairo, Egypt
- 12: Now at Ain Shams University, Cairo, Egypt
- 13: Also at National Centre for Nuclear Research, Swierk, Poland
- 14: Also at Université de Haute-Alsace, Mulhouse, France
- 15: Now at Joint Institute for Nuclear Research, Dubna, Russia
- 16: Also at Moscow State University, Moscow, Russia
- 17: Also at Brandenburg University of Technology, Cottbus, Germany
- 18: Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
- 19: Also at Eötvös Loránd University, Budapest, Hungary
- 20: Also at Tata Institute of Fundamental Research - HECR, Mumbai, India
- 21: Also at University of Visva-Bharati, Santiniketan, India
- 22: Also at Sharif University of Technology, Tehran, Iran
- 23: Also at Isfahan University of Technology, Isfahan, Iran
- 24: Also at Plasma Physics Research Center, Science and Research Branch, Islamic Azad University, Teheran, Iran
- 25: Also at Facoltà Ingegneria Università di Roma, Roma, Italy
- 26: Also at Università della Basilicata, Potenza, Italy
- 27: Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- 28: Also at Università degli Studi di Siena, Siena, Italy
- 29: Also at University of Bucharest, Faculty of Physics, Bucuresti-Magurele, Romania
- 30: Also at Faculty of Physics of University of Belgrade, Belgrade, Serbia
- 31: Also at University of California, Los Angeles, Los Angeles, USA
- 32: Also at Scuola Normale e Sezione dell' INFN, Pisa, Italy
- 33: Also at INFN Sezione di Roma; Università di Roma "La Sapienza", Roma, Italy
- 34: Also at University of Athens, Athens, Greece
- 35: Also at Rutherford Appleton Laboratory, Didcot, United Kingdom
- 36: Also at The University of Kansas, Lawrence, USA
- 37: Also at Paul Scherrer Institut, Villigen, Switzerland
- 38: Also at Institute for Theoretical and Experimental Physics, Moscow, Russia
- 39: Also at Gaziosmanpasa University, Tokat, Turkey
- 40: Also at Adiyaman University, Adiyaman, Turkey
- 41: Also at Izmir Institute of Technology, Izmir, Turkey
- 42: Also at The University of Iowa, Iowa City, USA
- 43: Also at Mersin University, Mersin, Turkey
- 44: Also at Ozyegin University, Istanbul, Turkey
- 45: Also at Kafkas University, Kars, Turkey
- 46: Also at Suleyman Demirel University, Isparta, Turkey
- 47: Also at Ege University, Izmir, Turkey
- 48: Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- 49: Also at INFN Sezione di Perugia; Università di Perugia, Perugia, Italy
- 50: Also at University of Sydney, Sydney, Australia
- 51: Also at Utah Valley University, Orem, USA

52: Also at Institute for Nuclear Research, Moscow, Russia

53: Also at University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia

54: Also at Argonne National Laboratory, Argonne, USA

55: Also at Erzincan University, Erzincan, Turkey

56: Also at KFKI Research Institute for Particle and Nuclear Physics, Budapest, Hungary

57: Also at Kyungpook National University, Daegu, Korea